

(1)  $\begin{pmatrix} 54 & 110 \\ -3 & -11 \end{pmatrix}$       (2)  $\frac{1}{66} \begin{pmatrix} -54 & -110 \\ 3 & 11 \end{pmatrix}$

(3)  $\frac{1}{66} \begin{pmatrix} 54 & 110 \\ 3 & 11 \end{pmatrix}$       (4) none of these

5. The value of the determinant  $\begin{vmatrix} \cos \alpha & \sin \alpha & 0 \\ -\sin \alpha & \cos \alpha & 0 \\ 0 & 0 & 1 \end{vmatrix}$  is :

- (1) -1      (2) 0      (3) 1      (4)  $\cos 2\alpha$

6. If  $\omega$  is the cube root of unity then the value of  $\begin{vmatrix} 1 & \omega & \omega^2 \\ \omega & 1 & \omega \\ \omega^2 & \omega & 1 \end{vmatrix}$  will be :

- (1)  $\omega$       (2)  $\omega^2 + 1$       (3) 0      (4) 1

7. If  $(1+x)^n = C_0 + C_1x + C_2x^2 + \dots + C_nx^n$ , then  $C_0 + \frac{C_1}{2} + \frac{C_2}{3} + \dots + \frac{C_n}{n+1}$  is equal to :

- (1)  $\frac{2^n - 1}{n + 1}$       (2)  $\frac{2^n - 1}{n - 1}$       (3)  $\frac{2^{n-1} - 1}{n + 1}$       (4)  $\frac{2^{n+1} - 1}{n + 1}$

8. If  $(1+x)^n = C_0 + C_1x + C_2x^2 + \dots + C_nx^n$ , then  $C_0C_2 + C_1C_3 + C_2C_4 + \dots = C_{n-2}C_n$  is equal to :

- (1)  $\frac{2n}{n-2} \frac{n-2}{n-2}$       (2)  $\frac{2n}{n} \frac{n}{n}$       (3)  $\frac{2n}{n-2} \frac{n+2}{n+2}$       (4) none of these

9. If the ratio of the second and third term in the expansion of  $(a+b)^2$  is equal to the ratio of third and fourth term in the expansion of  $(a+b)^{n+3}$  then the value of n is equal to :

- (1) 6      (2) 4      (3) 5      (4) 3

10. The number of different words that can be formed by using the letters of the word 'MISSISIPPI' is :

- (1) 5067      (2) 6705      (3) 1520      (4) 2520

11. If  $a^2, b^2, c^2$  are in A.P. then  $\frac{1}{b+c^2}, \frac{1}{c+a}, \frac{1}{a+b}$  will be :

- (1) in H.P.      (2) in G.P.      (3) in A.P.      (4) in arithmetic geometrico progression

12. A bag contains 5 white and 3 black balls. Two balls are drawn at random then the probability that out of the two one ball is red and other is black will be :

- (1)  $\frac{15}{56}$       (2)  $\frac{11}{28}$       (3)  $\frac{15}{28}$       (4)  $\frac{2}{7}$

13. Two dice are thrown together then the probability that the sum of numbers appearing on the dice is 7 :

- (1)  $\frac{1}{6}$       (2)  $\frac{1}{12}$       (3)  $\frac{5}{36}$       (4) none of these

14. The position vectors of the two points A and B are a and b respectively then the position vector of the point C which divides AB in the ration 2 : 1 will be :

- (1)  $\frac{1+b}{3}$       (2)  $\frac{2a+b}{3}$       (3)  $\frac{a+2b}{3}$       (4) none of these

15. If  $a + b = a - b$  , then the angle between a and b will be :

- (1)  $180^0$       (2)  $90^0$       (3)  $60^0$       (4)  $0^0$

16. The area of the region bounded by the curve  $y = \sin^2 x$  , x-axis and the lines  $x = 0$ ,  $x = \pi/2$  is

- (1)  $\pi$       (2)  $\pi/8$       (3)  $\frac{\pi}{4}$       (4)  $\frac{\pi}{2}$

17.  $\int_0^{\pi/2} \frac{\sin x}{\sqrt{1 + \sin 2x}} dx$  is equal to :

- (1)  $\pi$       (2)  $2\pi$       (3)  $\frac{\pi}{4}$       (4)  $\frac{\pi}{2}$

18.  $\int x e^x dx$  is equal to :

- (1)  $(x - 1)e^x + C$       (2)  $(1 - x)e^x + C$   
(3)  $(1 - x)e^x + C$       (4) none of these

19.  $\int x^2 \sin x^3 dx$  is equal to :

- (1)  $\frac{1}{3} \sin x^3 + C$       (2)  $-\frac{1}{3} \sin x^3 + C$   
(3)  $\frac{1}{3} \cos x^3 + C$       (4)  $-\frac{1}{3} \cos x^3 + C$

20. The max. value of  $\sin x + \cos x$  :

- (1) 1      (2)  $\frac{1}{\sqrt{2}}$       (3)  $\sqrt{2}$       (4) none of these

21. The angle between the curves  $y = x$  and  $y^2 = 4x$  at origin will be :

- (1) 0                      (2)  $\frac{\pi}{3}$                       (3)  $\frac{\pi}{4}$                       (4) none of these

**22. If the volume of a balloon is increasing at the rate of 25 cm<sup>3</sup>/sec., then if the radius of the balloon is 5 cm. then the rate of change of the surface area is :**

- (1) 20 cm<sup>2</sup>/sec.                      (2) 10 cm<sup>2</sup>/sec.                      (3) 5 cm<sup>2</sup>/sec.                      (4) 10 cm./sec.

**23. The differential coefficient of x<sup>x</sup> is :**

- (1) x<sup>x</sup> log<sub>e</sub> x                      (2) x<sup>x</sup> (1 + log<sub>e</sub> x)                      (3) x<sup>x</sup> (1 - log<sub>e</sub> x)                      (4) none of these

**24.  $\frac{d}{dx} (\sin x)^{\tan x}$  is equal to :**

- (1)  $(\sin x)^{\tan x} [1 - \sec^2 x \log \sin x]$   
 (2)  $(\tan x)^{\sin x} \log \sec^2 x \log \sin x]$   
 (3)  $(\sin x)^{\tan x} [1 + \sec^2 x \log \sin x]$   
 (4) none of these

**25. If  $y = \sec^{-1} \left( \frac{x+1}{x-1} \right) + \sin^{-1} \left( \frac{x-1}{x+1} \right)$**

**then  $\frac{dy}{dx}$  is equal to :**

- (1)  $\infty$                       (2) 0                      (3) 1                      (4) -1

**26. If  $y = \sin^{-1} \left( \frac{2x}{1+x^2} \right)$ , then  $\frac{dy}{dx}$  is equal to :**

- (1)  $-\frac{1}{1+x^2}$                       (2)  $-\frac{2}{1+x^2}$   
 (3)  $\frac{2}{1+x^2}$                       (4)  $\frac{2x}{1+x^2}$

**27. If  $x\sqrt{1+y} + y\sqrt{1+x} = 0$  then  $\frac{dy}{dx}$  is equal to :**

- (1)  $\frac{1}{1+x^2}$                       (2)  $-\frac{1}{1+x^2}$   
 (3)  $\frac{2}{(1+x)^2}$                       (4) none of these

**28. The continuous product of the roots of (-1)<sup>2/3</sup> is :**

- (1)  $\omega^2$                       (2)  $\omega$                       (3) 0                      (4) 1

**29. The value of  $\sin^{-1} x$  :**

- (1)  $\log (x - \sqrt{x^2 - 1})$                       (2)  $\log (x + \sqrt{x^2 - 1})$

(3)  $\log(x + \sqrt{x^2 + 1})$

(4)  $\frac{1}{2} \log \frac{1+x}{1-x}$

30. The equation of a st-line passing through the point (1,2) and making equal angles to with axes, will be :

(1)  $x - y - 2 = 0$

(2)  $x + y + 1 = 0$

(3)  $x - y = 1$

(4)  $x + y = 1$

31. If the vertices of a parallelogram are (0,0), (2,1), (1,3) and (1,2) then the angle between their diagonals will be:

(1)  $\frac{\pi}{4}$

(2)  $\frac{3\pi}{2}$

(3)  $\frac{\pi}{2}$

(4)  $\frac{\pi}{3}$

32. The equation of line which is parallel to the straight line  $3x + 4y - 7 = 0$  and passing through (1,2) is :

(1)  $3x + 4y = 11$

(2)  $3x + 4y + 11 = 0$

(3)  $4x - 3y + 2 = 0$

(4)  $3x + 4y + 7 = 0$

33. The pole of the straight line  $9x + y - 28 = 0$  w.r.t. the circle  $x^2 + y^2 = 16$  will be:

(1)  $\left(\frac{33}{7}, \frac{3}{7}\right)$

(2)  $\left(\frac{33}{7}, \frac{4}{7}\right)$

(3)  $\left(\frac{4}{7}, \frac{36}{7}\right)$

(4)  $\left(\frac{36}{7}, \frac{4}{7}\right)$

34. The equation of the tangent from origin to the circle  $x^2 + y^2 - 2rx - 2hy + h^2 = 0$  is:

(1)  $(h^2 - r^2)x + 2rhy = 0$

(2)  $y = 0$

(3)  $x - y = 0$

(4)  $(h^2 - r^2)x - 2rhy = 0$

35. If a tangent at a point p to the parabola meets to the directrix at Q. If S is the focus of the parabola then  $\angle PSQ$  is equal to :

(1)  $\pi$

(2)  $\frac{\pi}{2}$

(3)  $\frac{\pi}{3}$

(4)  $\frac{\pi}{4}$

36. If  $f(y) = \log y$ , then  $f(y) + f(1/y)$  is equal to :

(1) 2

(2) 0

(3) -1

(4) 1

37.  $\lim_{x \rightarrow 0^+} \frac{e^x - \log(1+x)}{x^2}$  is equal to :

(1)  $\frac{1}{2}$

(2)  $\frac{1}{3}$

(3)  $\frac{3}{2}$

(4)  $\frac{2}{3}$

38. If  $\alpha$  and  $\beta$  are the roots of the equation  $1 - (1 + n^2 + n^4) = 0$  then  $\alpha + \beta$  is equal to :

- (1)  $2n^2$       (2)  $n^2$       (3)  $-n^2$       (4)  $n^2 + 2$

39. The H.M. between 1 and  $\frac{1}{16}$  will be :

- (1)  $\frac{17}{2}$       (2)  $\frac{2}{17}$       (3)  $\frac{17}{32}$       (4)  $\frac{32}{17}$

40. If for two numbers G.M. is 4 and A.M. is 5, then H.M. will be :

- (1)  $\frac{25}{15}$       (2)  $\frac{17}{8}$       (3)  $\frac{16}{5}$       (4)  $\frac{5}{16}$

41. If  ${}^{10}C_r = {}^{10}C_{r+2}$  then  ${}^5C_r$  is equal to :

- (1) 360      (2) 120      (3) 10      (4) 5

42. The value of  $1 + \frac{1}{4} + \frac{1 \cdot 3}{4 \cdot 8} + \frac{1 \cdot 3 \cdot 5}{4 \cdot 8 \cdot 12} + \dots$  is :

- (1)  $\sqrt{3/2}$       (2)  $\sqrt{2}$       (3) 2      (4)  $3/2$

43. If  $(1+x)^n = C_0 + C_1x + C_2x^2 + \dots + C_nx^n$  then  $\frac{C_1}{C_0} + \frac{2C_2}{C_1} + \frac{3C_3}{C_2} + \dots + \frac{nC_n}{C_{n-1}}$  is equal to :

- (1)  $\frac{n(n+1)}{2}$       (2)  $\frac{n(n^2+1)}{2}$       (3)  $\frac{n(n+1)}{n!}$       (4)  $\frac{n(n-1)}{2}$

44. In the expansion of  $\left(2^4 - \frac{1}{x^7}\right)$  the term independent of x is :

- (1) -32190      (2) 114050      (3) 42240      (4) 330

45. The value of the determinant  $\begin{vmatrix} 4 & -6 & 1 \\ -1 & -1 & 1 \\ -4 & 11 & -1 \end{vmatrix}$  is :

- (1) 0      (2) -25      (3) 25      (4) none of these

46. If  $\begin{vmatrix} 1 & 2 & 4 \\ 3 & 6+x & 7 \end{vmatrix} = 0$ , then the value of x will be :

5 10 4+x

- (1) 3                      (2) 0                      (3) 1                      (4) none of these


47. If  $A = \begin{pmatrix} d_1 & 0 & 0 \\ 0 & d_2 & 0 \\ 0 & 0 & d_3 \end{pmatrix}$  then  $\text{adj } A =$

(1)  $\begin{pmatrix} d_1^{-1} & 0 & 0 \\ 0 & d_2^{-1} & 0 \\ 0 & 0 & d_3^{-1} \end{pmatrix}$

(2)  $\begin{pmatrix} d_2 d_3 & 0 & 0 \\ 0 & d_1 d_3 & 0 \\ 0 & 0 & d_1 d_2 \end{pmatrix}$

(3)  $\begin{pmatrix} d_2 d_2 & 0 & 0 \\ 0 & d_1 d_3 & 0 \\ 0 & 0 & d_1 d_3 \end{pmatrix}$

(4)  $\begin{pmatrix} d_1 d_3 & 0 & 0 \\ 0 & d_2 d_3 & 0 \\ 0 & 0 & d_1 d_2 \end{pmatrix}$



48. If  $A = \begin{pmatrix} 2 & 4 \\ 0 & 3 \end{pmatrix}$  and  $B = \begin{pmatrix} 1 & 2 \\ 0 & 5 \end{pmatrix}$ , then  $4A - 3B$  is equal to :

- (1)  $\begin{pmatrix} 1 & 2 \\ 0 & 2 \end{pmatrix}$                       (2)  $\begin{pmatrix} -5 & -10 \\ 0 & 3 \end{pmatrix}$                       (3)  $B = \begin{pmatrix} 5 & 10 \\ 0 & -3 \end{pmatrix}$                       (4)  $A = \begin{pmatrix} 7 & 14 \\ 0 & 7 \end{pmatrix}$

49. If  $A = \begin{pmatrix} \cos x & \sin x \\ -\sin x & \cos x \end{pmatrix}$ , then  $A^{-1}$  is equal to :

- (1)  $\begin{pmatrix} \cos x & \sin x \\ \sin x & \cos x \end{pmatrix}$  (2)  $\begin{pmatrix} \cos x & -\sin x \\ \sin x & \cos x \end{pmatrix}$  (3)  $\begin{pmatrix} \cos x & \sin x \\ -\sin x & \cos x \end{pmatrix}$  (4) none of these

50. A card is drawn at random from a pack of playing cards. The probability that it is red or an ace, is :

- (1)  $\frac{1}{13}$  (2)  $\frac{1}{52}$  (3)  $\frac{17}{52}$  (4)  $\frac{4}{13}$

51. If the sum of two unit vector is also a unit vector then the magnitude of their difference will be :

- (1) 1 (2)  $\sqrt{3}$  (3)  $\frac{1}{\sqrt{3}}$  (4)  $\sqrt{2}$

52. The unit vector perpendicular to the vectors  $6\mathbf{i} + 2\mathbf{j} + 3\mathbf{k}$  and  $3\mathbf{i} - 6\mathbf{j} - 2\mathbf{k}$  will be:

- (1)  $\frac{2\mathbf{i} - 3\mathbf{j} - 6\mathbf{k}}{7}$  (2)  $\frac{2\mathbf{i} - 3\mathbf{j} + 6\mathbf{k}}{7}$  (3)  $\frac{2\mathbf{i} + 3\mathbf{j} - 6\mathbf{k}}{7}$  (4)  $\frac{2\mathbf{i} + 3\mathbf{j} + 6\mathbf{k}}{7}$

53. The area of the region bounded by the curves  $y^2 = 4ax$ ,  $x = 0$  and  $x = a$  is

- (1)  $4\pi a^2$  (2)  $3\pi a^2$  (3)  $2\pi a^2$  (4)  $\pi a^2$

54. The area of the region bounded by the curves  $y^2 = 4ax$ ,  $x = 0$  and  $x = a$  is :

- (1)  $\frac{5}{3} a^2$  (2)  $\frac{2}{3} a^2$  (3)  $\frac{8}{3} a^2$  (4)  $\frac{4}{3} a^2$

55.  $\int \cos^3 x \, dx$  is equal to

- (1)  $\frac{\sin 3x}{4} + 3 \sin x + C$   
 (2)  $\frac{\sin 3x}{3} + \frac{\sin x}{2} + C$   
 (3)  $\sin 3x + C$   
 (4)  $\frac{\sin 3x}{12} + \frac{3}{4} \sin x + C$

56. If  $x = a(t + \sin t)$  and  $y = a(1 - \cos t)$  then  $\frac{dy}{dx}$  is equal to :

- (1)  $\tan t$  (2)  $\tan 2t$  (3)  $\cot(t/2)$  (4)  $\tan(t/2)$

57. If  $x = t^2$  and  $y = 2t$ , then the normal at  $t = 1$  is :

- (1)  $x + y - 3 = 0$       (2)  $x + y - 1 = 0$   
 (3)  $x + y + 1 = 0$       (4)  $x + y + 3 = 0$

**58.  $f(x) = 2x^3 - 9x^2 + 12x + 29$  is a monotonic decreasing function when :**

- (1)  $1 < x < 2$     (2)  $x > 1$       (3)  $x > 2$       (4)  $x < 2$

**59. The height of the cylinder of maximum volume that can be inscribed in a sphere of radius  $r$  is :**

- (1)  $2\sqrt{3}r$       (2)  $\frac{2r}{\sqrt{3}}$       (3)  $r\sqrt{3}$       (4)  $\frac{r}{\sqrt{3}}$

**60.  $\int \sec x \, dx$  is equal to :**

- (1)  $\log \sin x + C$   
 (2)  $\log \tan (x/2) + C$   
 (3)  $-\log (\sec x - \tan x) + C$   
 (4)  $\log \tan \left[ \frac{\pi}{2} + \frac{\pi}{4} \right] + C$

**61. The differential coefficient of  $\sin^{-1} \left( \frac{1-x^2}{1+x^2} \right)$  w.r.t.  $x$  is :**

- (1)  $-\frac{2}{1+x^2}$     (2)  $\frac{2}{1+x^2}$       (3)  $\frac{1}{1+x^2}$     (4) none of these

**62.  $d(\sec^{-1} x)$  is equal to :**

- (1)  $\frac{1}{x\sqrt{x-1}}$       (2)  $\frac{1}{x\sqrt{x^2-1}}$       (3)  $\frac{1}{x\sqrt{1+x^2}}$       (4)  $\frac{1}{x\sqrt{1-x^2}}$

**63. The differential coefficient of  $\tan^{-1} \frac{1-x^2}{\sqrt{1+x^2}}$  w.r.t.  $x$  is :**

- (1)  $\frac{1}{2}$       (2)  $1$       (3)  $-\frac{1}{2}$       (4) none of these

**64.  $\lim_{x \rightarrow 0} \frac{\tan 2x - x}{3x - \sin x}$  is equal to :**

- (1)  $0$       (2)  $1$       (3)  $\frac{1}{3}$       (4)  $\frac{1}{4}$

**65. The differential coefficient of  $\sin^{-1} x$  w.r.t.  $\cos^{-1} \sqrt{1-x^2}$  is :**

- (1)  $-\frac{1}{\sqrt{1+x^2}}$       (2)  $\frac{1}{\sqrt{1-x^2}}$       (3)  $\frac{2}{\sqrt{1-x^3}}$       (4) none of these

**66. The sum of 20 terms of the series  $1 + 4 + 7 + 10 + \dots$  is :**



- (1) 290      (2) 490      (3) 590      (4) none of these

67. Which terms of the series  $\frac{1}{2}, \frac{-1}{2}, 1-2, \dots$  is  $-128$  :

- (1) 10<sup>th</sup>      (2) 8<sup>th</sup>      (3) 9<sup>th</sup>      (4) 12<sup>th</sup>

68. If  ${}^n P_4 : {}^n P_5 = 1 : 2$ , then  $n$  is equal to :

- (1) 2      (2) 4      (3) 5      (4) 6

69.  $\frac{(\cos 3\theta - i \sin 3\theta)\theta^5 (\cos 2\theta + i \sin 2\theta)\theta^4}{(\cos \theta + i \sin \theta)\theta^4 (\cos \theta - i \sin \theta)\theta^6}$  is equal to :

- (1)  $\cos 27\theta - i \sin 27\theta$   
(2)  $\cos 33\theta - i \sin 33\theta$   
(3)  $\cos 33\theta + i \sin 33\theta$   
(4)  $\cos 27\theta + i \sin 27\theta$

70. The value of  $\cos^{-1} x$  is :

- (1)  $\log(x - \sqrt{x^2 - 1})$       (2)  $\log(x + \sqrt{x^2 - 1})$   
(3)  $\log(x + \sqrt{x^2 + 1})$       (4)  $\log(x - \sqrt{x^2 + 1})$

71. Find the equation of the straight line which is perpendicular to the line  $\frac{x}{a} - \frac{y}{b} = 1$  and passes through the point where the given st-line cuts the x-axis :

- (1)  $ax + by = a^2$       (2)  $ax - by = a^2$       (3)  $ax + by = b^2$       (4)  $bx - ay = ab$

72. If the lines  $x + y = 1$ ,  $2x - y = 0$  and  $x + 2y + \lambda = 0$  are concurrent then  $\lambda$  is equal to :

- (1)  $-\frac{2}{3}$       (2)  $\frac{2}{3}$       (3)  $-\frac{5}{3}$       (4)  $\frac{5}{3}$

73. If two vertices of a triangle are (6,4), (2,6) and its centroid is (4, 6) then its third vertex will be :

- (1) (6,4)      (2) (8,4)      (3) (4,8)      (4) none of these

74. The radical axis of the circles  $2x^2 + 2y^2 - 7x = 0$  and  $x^2 + y^2 - 4y - 7 = 0$  is :

- (1)  $8x - 7y + 14 = 0$       (2)  $7x - 8y + 14 = 0$   
(3)  $7x - 8y - 14 = 0$       (4)  $7x + 8y + 14 = 0$

75. The equation of the polar line w.r.t. the pole (1, -2) to the circle  $x^2 + y^2 - 2x - 6y + 5 = 0$  is :

- (1)  $x + y - 1 = 0$       (2)  $x + y + 1 = 0$       (3)  $y = 2$       (4)  $x = 2$

76. The vertex of the parabola  $x^2 - y + 6x + 10 = 0$  is :

- (1) (3,1)                      (2) (3, -1)                      (3) (3,-2)                      (4) (-3, 1)

77. If  $f(\theta) = \tan \theta$ , then the value of  $\frac{f(\theta) - f(\phi)}{1 + f(\theta)f(\phi)}$  is :

- (1)  $\theta - \phi$                       (2)  $f(\theta / \phi)$                       (3)  $f(\theta - \phi)$                       (4)  $f(\theta + \phi)$

78.  $\lim_{x \rightarrow 0} \frac{x^2 - 3x + 2}{2x^2 + x - 3}$  is equal to :

- (1) 0                      (2) 2                      (3)  $\frac{1}{2}$                       (4)  $\infty$

79.  $\lim_{x \rightarrow 0} \left( \frac{\sqrt{1+x} - \sqrt{1-x}}{x} \right)$  is equal to :

- (1) -1                      (2) 1                      (3) 2                      (4)  $\frac{1}{2}$

80. The equation of the normal at a point of intersection of line  $2x + y = 3$  and curve  $yx^2 + y^2 = 5$  is :

- (1)  $2x + 2y + 3 = 0$                       (2)  $x - y + 4 = 0$                       (3)  $x - 4y + 3 = 0$                       (4)  $x + y + 2 = 0$

81. If  $f(x) = \frac{x - 3}{x + 1}$ , then  $f[f(f(x))]$  is equal to :

- (1)  $\frac{-1}{x}$                       (2)  $-x$                       (3)  $\frac{1}{x}$                       (4)  $x$

82. The modulus of  $\frac{1+i}{1-i}$  is :

- (1)  $\sqrt{2}$                       (2) 2                      (3)  $\frac{1}{2}$                       (4) 1

83. The value of  $\frac{4\sqrt{3}-3}{7} - \frac{\sqrt{3}-3}{7}$  is

- (1)  $\frac{3\sqrt{3}}{7}$                       (2)  $-\frac{3\sqrt{3}}{7}$                       (3)  $\frac{3\sqrt{3}}{7} i$                       (4) none of these

84.  $\frac{1-2i}{2+i} + \frac{4-i}{+2i}$  is equal to :

- (1)  $\frac{10}{13} + \frac{24}{13} i$                       (2)  $\frac{10}{13} - \frac{24}{13} i$                       (3)  $\frac{24}{13} - \frac{10}{13} i$                       (4)  $\frac{24}{13} + \frac{10}{13} i$

85. If  $z = 5 + 3i$  then the value of  $|z - 2|$  will be :

- (1)  $\sqrt{13}$                       (2)  $2\sqrt{3}$                       (3)  $3\sqrt{2}$                       (4) 13

86. The imaginary part of  $\frac{1-i}{1+i}$  is :

- (1)  $-i$       (2)  $-1$       (3)  $1$       (4)  $i$

87. If  $z_1 = 1 + 2i$  and  $z_2 = i - 1$ , then  $z_1/z_2$  is equal to :

- (1)  $\frac{1}{2} - \frac{3}{2}i$       (2)  $-\frac{1}{2} + \frac{3}{2}i$       (3)  $\frac{1}{2} - \frac{3}{2}i$       (4) none of these

88. The amplitude of  $1 - \sqrt{3}i$  is :

- (1)  $-\frac{2\pi}{3}$       (2)  $-\frac{\pi}{3}$       (3)  $\frac{2\pi}{3}$       (4)  $\frac{\pi}{3}$

89. If  $\alpha$  and  $\beta$  are the roots of the equation  $x^2 + px + q = 0$  then the value of  $\alpha^3 + \beta^3 = 0$  will be :

- (1)  $p^3 - 3pq$       (2)  $-(p^3 + 3pq)$       (3)  $p^3 + 3pq$       (4)  $-p^3 + 3pq$

90. If  $\alpha$  and  $\beta$  are the roots of the equation whose roots are  $\frac{1}{\alpha}, \frac{1}{\beta}$  is :

- (1)  $x^2 + x + 1 = 0$       (2)  $x^2 - x + 1 = 0$       (3)  $x^2 - x = 1$       (4)  $x^2 - x = 1$

91. If  $z = \frac{(1+i)(2+i)}{3+i}$  then  $|z|$  is equal to :

- (1)  $-\frac{1}{2}$       (2)  $\frac{1}{2}$       (3)  $1$       (4)  $-1$

92. The slope of the tangent to the parabola  $y^2 = 4ax$  at point  $(at^2, 2at)$  will be :

- (1)  $-t$       (2)  $-1/t$       (3)  $1/t$       (4)  $t$

93. If  $a = i - 2j$  and  $b = 2i + \lambda j$  are the parallel vectors then  $\lambda$  is equal to :

- (1)  $-2$       (2)  $2$       (3)  $-4$       (4)  $4$

94. A stone is thrown in silent water, the ripples are moving at the rate of  $6 \text{ cm/sec}$ . then the rate of change of the are when the radius of the unite is  $10 \text{ cm}$ . at the time when radius of the circle is  $10 \text{ cm}$ , then the rate at which its area increases is :

- (1)  $120 \text{ m}^2/\text{sec}$ .      (2)  $\pi \text{ cm}^2/\text{sec}$ .      (3)  $120 \text{ cm}^2/\text{sec}$ .      (4)  $120 \pi \text{ cm}^2/\text{sec}$ .

95. A dice is thrown then the probability that the sum of the number is  $1$  or  $6$  is :

- (1)  $1/6$       (2)  $1/3$       (3)  $2/3$       (4)  $3/4$

96. The value of  $\cos h(\pi i)$  is :

- (1)  $0$       (2)  $1$       (3)  $-1$       (4) none of these

97. For  $Z_1, Z_2 \in \mathbb{C}$  the value of  $|Z_1 + Z_2|^2 + |Z_1 - Z_2|^2$  will be :

- (1)  $2(|Z_1|^2 + |Z_2|^2)$       (2)  $|Z_1|^2 + |Z_2|^2$       (3)  $|Z_1|^2 + |Z_2|^2 - |Z_1| - |Z_2|$       (4)  $2(|Z_1|^2 - |Z_2|^2)$

