

## Section 4: Mathematics / Biology

Students will have to attempt either Mathematics/Biology as per the eligibility of the program applied.

### Mathematics

66. The solution of the equation.  $\log\left(\log_5\left(\sqrt{x+5} + \sqrt{x}\right)\right) = 0$  is
- (a) 2                      (b) 4                      (c) 3                      (d) 8
67. Let  $\frac{1}{q+r}$ ,  $\frac{1}{r+p}$  and  $\frac{1}{p+q}$  are in A.P. where  $p, q, r, \neq 0$ , then
- (a)  $p, q, r$  are in A.P.                      (b)  $p^2, q^2, r^2$  are in A.P.  
(c)  $\frac{1}{p} \cdot \frac{1}{q} \cdot \frac{1}{r}$  in A.P.                      (d) none of these
68. If  $b \in \mathbb{R}^+$  then the roots of the equation  $(2+b)x^2 + (3+b)x + (4+b) = 0$  is
- (a) real and distinct      (b) real and equal      (c) imaginary      (d) cannot predicted
69. Solve for integral solutions  $x_1 + x_2 + x_3 + \dots + x_6 \leq 17$ , where  $1 \leq x_i \leq 6, i = 1, 2, \dots, 6$ .  
Number of solutions will be
- (a)  ${}^{17}C_6 - 6{}^{11}C_5$       (b)  ${}^{17}C_{11} - 6{}^{11}C_5$       (c)  ${}^{17}C_5 - 6{}^{11}C_5$       (d)  ${}^{17}C_{11} - 5{}^{11}C_6$
70. The probability that a certain beginner at golf gets a good shot if he uses the correct club is  $\frac{1}{3}$ , and the probability of a good shot with an incorrect club is  $\frac{1}{4}$ . In his bag there are 5 different clubs, only one of which is correct for the shot in question. If he chooses a club at random and take a stroke, the probability that he gets a good shot is
- (a)  $\frac{1}{3}$                       (b)  $\frac{1}{12}$                       (c)  $\frac{4}{15}$                       (d)  $\frac{7}{12}$

71. OPQR is a square and M, N are the middle points of the side PQ and QR respectively. Then the ratio of the area of the square and the triangle OMN is

- (a) 4 : 1                      (b) 2 : 1                      (c) 4 : 3                      (d) 8 : 3

72. Two vertices of an equilateral triangle are  $(-1, 0)$  and  $(1, 0)$  and its circumcircle is

- (a)  $x^2 + \left(y - \frac{1}{\sqrt{3}}\right)^2 = \frac{4}{3}$                       (b)  $x^2 - \left(y + \frac{1}{\sqrt{3}}\right)^2 = \frac{4}{3}$   
(c)  $x^2 + \left(y - \frac{1}{\sqrt{3}}\right)^2 = -\frac{4}{3}$                       (d) none of these

73. If in a  $\Delta ABC$ ,  $\sin^2 A + \sin^2 B + \sin^2 C = 2$ , then the triangle is always

- (a) isosceles triangle                      (b) right angled                      (c) acute angled                      (d) obtuse angled

74. If the vertex and the focus of a parabola are  $(-1, 1)$  and  $(2, 3)$  respectively, then the equation of the directrix is

- (a)  $3x + 2y - 25 = 0$                       (b)  $x + 2y + 7 = 0$                       (c)  $2x - 3y + 10 = 0$                       (d)  $3x + 2y + 14 = 0$ .

75. The radius of the circle passing through the foci of the ellipse  $\frac{x^2}{16} + \frac{y^2}{9} = 1$  and having its centre at  $(0, 3)$  is

- (a) 4                      (b) 3                      (c)  $\sqrt{12}$                       (d)  $7/2$

76. If  $P(x_1, y_1)$ ,  $Q(x_2, y_2)$ ,  $R(x_3, y_3)$  and  $S(x_4, y_4)$  are four concyclic points on the rectangular hyperbola  $xy = c^2$ , then the co-ordinates of the orthocentre of  $\Delta PQR$  are

- (a)  $(x_4, -y_4)$                       (b)  $(x_4, y_4)$                       (c)  $(-x_4, -y_4)$                       (d)  $(-x_4, y_4)$

77. The coefficient of  $x^n y^n$  in the expansion of  $[(1 + x)(1 + y)(x + y)]^n$  is

- (a)  $\sum_{r=0}^n C_r$                       (b)  $\sum_{r=0}^n C_r^2$                       (c)  $\sum_{r=0}^n C_r^3$                       (d) none of these

78.  $z_0$  is one of the roots of the equation  $z^n \cos \theta_0 + z^{n-1} \cos \theta_1 + \dots + \cos \theta_n = 2$ , where  $\theta_i \in \mathbb{R}$ , then

- (a)  $|z_0| < \frac{1}{2}$       (b)  $|z_0| > \frac{1}{2}$       (c)  $|z_0| = \frac{1}{2}$       (d) none of these

79. The second order differential equation is

- (a)  $y'^2 + x + y^2$       (b)  $y'y'' + y = \sin x$       (c)  $y'''' + y'' + y = 0$       (d)  $y' = 0$

80.  $\int e^{3x} \left( \frac{1+3\sin x}{1+\cos x} \right) dx$  is equal to

- (a)  $e^{3x} \cot x + c$       (b)  $e^{3x} \tan \frac{x}{2} + c$       (c)  $e^{3x} \sin x + c$       (d)  $e^{3x} \cos x + c$

81. If  $m$  and  $n$  are positive integers and  $f(x) = \int_1^x (t-a)^{2n} (t-b)^{2m+1} dt$ ,  $a \neq b$ , then

- (a)  $x = b$  is a point of local minimum      (b)  $x = b$  is a point of local maximum  
 (c)  $x = a$  is a point of local minimum      (d)  $x = a$  is a point of local maximum

82. If in a triangle ABC  $\frac{2\cos A}{a} + \frac{\cos B}{b} + \frac{2\cos C}{c} = \frac{a}{bc} + \frac{b}{ca}$ , then the value of the angle A is

- (a)  $45^\circ$       (b)  $90^\circ$       (c)  $135^\circ$       (d)  $60^\circ$

83. The general solution of the equation  $2^{\cos 2x} + 1 = 3 \cdot 2^{-\sin^2 x}$  is

- (a)  $n\pi$       (b)  $\left(n + \frac{1}{2}\right)\pi$       (c)  $\left(n - \frac{1}{2}\right)\pi$       (d) all of the above.

84. Total number of positive real values of  $x$  satisfying  $2[x] = x + \{x\}$ , where  $[.]$  and  $\{.\}$  denote the greatest integer function and fractional part respectively is equal to

- (a) 2      (b) 1      (c) 0      (d) 3

85. If  $\lim_{x \rightarrow 0} \frac{((a-n)nx - \tan x) \sin nx}{x^2} = 0$ , where  $n$  is nonzero real number, then  $a$  is equal to

- (a) 0      (b)  $\frac{n+1}{n}$       (c)  $n$       (d)  $n + \frac{1}{n}$

86.  $f(x) = \begin{cases} 4x - x^3 + \ln(a^2 - 3a + 3), & 0 \leq x < 3 \\ x - 18, & x \geq 3 \end{cases}$ . Find the complete set of values of  $a$  such that

$f(x)$  has a local minima at  $x = 3$  is

- (a)  $[-1, 2]$       (b)  $(-\infty, 1) \cup (2, \infty)$       (c)  $[1, 2]$       (d)  $(-\infty, -1) \cup (2, \infty)$

87. The number of values of  $k$  for the system of equations  $(k + 1)x + 8y = 4k$  and  $kx + (k + 3)y = 3k - 1$  has infinitely many solutions

- (a) 0                      (b) 1                      (c) 2                      (d) infinite

88. The matrix  $\begin{bmatrix} \frac{1+i}{2} & \frac{-1+i}{2} \\ \frac{1+i}{2} & \frac{1-i}{2} \end{bmatrix}$  is

- (a) unitary                      (b) null matrix                      (c) symmetric                      (d) none of these

89. The area between the curves  $y = xe^x$  and  $y = xe^{-x}$  and the line  $x = 1$  is

- (a)  $2e$                       (b)  $e$                       (c)  $2/e$                       (d)  $1/e$

90. If the unit vectors  $\vec{a}$  and  $\vec{b}$  are inclined at an angle  $2\theta$  and  $|\vec{a} - \vec{b}| < 1$  then  $\theta$  lies in the interval

- (a)  $\left[0, \frac{\pi}{6}\right)$                       (b)  $\left[\frac{5\pi}{6}, 2\pi\right]$                       (c)  $\left[\frac{\pi}{6}, \frac{\pi}{2}\right]$                       (d)  $\left[\frac{\pi}{2}, \frac{5\pi}{6}\right]$

