

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

1. Determine whether each of the following relations are reflexive, symmetric and transitive:

(i) Relation R in the set $A = \{1, 2, 3, \dots, 13, 14\}$ defined as

$$R = \{(x, y) : 3x - y = 0\}$$

(ii) Relation R in the set N of natural numbers defined as

$$R = \{(x, y) : y = x + 5 \text{ and } x < 4\}$$

(iii) Relation R in the set $A = \{1, 2, 3, 4, 5, 6\}$ as

$$R = \{(x, y) : y \text{ is divisible by } x\}$$

(iv) Relation R in the set Z of all integers defined as

$$R = \{(x, y) : x - y \text{ is an integer}\}$$

(v) Relation R in the set A of human beings in a town at a particular time given by

(a) $R = \{(x, y) : x \text{ and } y \text{ work at the same place}\}$

(b) $R = \{(x, y) : x \text{ and } y \text{ live in the same locality}\}$

(c) $R = \{(x, y) : x \text{ is exactly 7 cm taller than } y\}$

(d) $R = \{(x, y) : x \text{ is wife of } y\}$

(e) $R = \{(x, y) : x \text{ is father of } y\}$

Solution:

$$(i) R = \{(x, y) : 3x - y = 0\}$$

$$A = \{1, 2, 3, 4, 5, 6, \dots, 13, 14\}$$

Therefore, $R = \{(1, 3), (2, 6), (3, 9), (4, 12)\} \dots (1)$

As per reflexive property: (x, x)

$\in R$, then R is reflexive) Since there is no such pair, so R is not reflexive.

As per symmetric property: (x, y)

$\in R$ and $(y, x) \in R$, then R is symmetric. Since there is no such pair, R is not symmetric

As per transitive property: If (x, y)

$\in R$ and $(y, z) \in R$, then $(x, z) \in R$. Thus R is transitive. From (1), $(1, 3)$

$\in R$ and $(3, 9) \in R$ but $(1, 9) \notin R$, R is not transitive. Therefore, R is neither reflexive, nor symmetric and no

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

(ii) $R = \{(x, y) : y = x + 5 \text{ and } x < 4\}$ in set N of natural numbers.

Values of x are 1, 2, and 3

So, $R = \{(1, 6), (2, 7), (3, 8)\}$

As per reflexive property: (x, x)

$\in R$, then R is reflexive. Since there is no such pair, R is not reflexive.

As per symmetric property: (x, y)

$\in R$ and $(y, x) \in R$, then R is symmetric. Since there is no such pair, so R is not symmetric.

As per transitive property: If (x, y)

$\in R$ and $(y, z) \in R$, then $(x, z) \in R$. Thus R is transitive. Since there is no such pair, so R is not transitive.

Therefore, R is neither reflexive, nor symmetric and nor transitive.

(iii) $R = \{(x, y) : y \text{ is divisible by } x\}$ in $A = \{1, 2, 3, 4, 5, 6\}$

From above we have,

$R = \{(1, 1), (1, 2), (1, 3), (1, 4), (1, 5), (1, 6), (2, 2), (2, 4), (2, 6), (3, 3), (3, 6), (4, 4), (5, 5), (6, 6)\}$

As per reflexive property: (x, x)

$\in R$, then R is reflexive. $(1, 1), (2, 2), (3, 3), (4, 4), (5, 5)$ and $(6, 6)$

$\in R$. Therefore, R is reflexive. As per symmetric property: (x, y)

$\in R$ and $(y, x) \in R$, then R is symmetric. $(1, 2)$

$\in R$ but $(2, 1) \notin R$. So R is not symmetric. As per transitive property: If (x, y)

$\in R$ and $(y, z) \in R$, then $(x, z) \in R$. Thus R is transitive. Also $(1, 4)$

$\in R$ and $(4, 4) \in R$ and $(1, 4) \in R$, So R is transitive. Therefore, R is reflexive and transitive but not symmetric.

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

(iv) $R = \{(x, y) : x - y \text{ is an integer}\}$ in set Z of all integers.

Now, (x, x) , say $(1, 1) = x - y = 1 - 1 = 0$

$\in Z \Rightarrow R$ is reflexive. (x, y)

$\in R$ and $(y, x) \in R$, i.e., $x - y$ and $y - x$ are integers $\Rightarrow R$ is symmetric.

(x, y)

$\in R$ and $(y, z) \in R$, then $(x, z) \in R$ i.e., $x - y$ and $y - z$ and $x - z$ are integers.

(x, z)

$\in R \Rightarrow R$ is transitive. Therefore, R is reflexive, symmetric and transitive.

(v)

(a) $R = \{(x, y) : x \text{ and } y \text{ work at the same place}\}$

For reflexive: x and x can work at same place

(x, x)

$\in R$ R is reflexive.

For symmetric: x and y work at same place so y and x also work at same place.

(x, y)

$\in R$ and $(y, x) \in R$ R is symmetric.

For transitive: x and y work at same place and y and z work at same place, then x and z also work at same place.

(x, y)

$\in R$ and $(y, z) \in R$ then $(x, z) \in R$ R is transitive

Therefore, R is reflexive, symmetric and transitive.

(b) $R = \{(x, y) : x \text{ and } y \text{ live in the same locality}\}$

(x, x)

$\in R \Rightarrow R$ is reflexive. (x, y)

$\in R$ and $(y, x) \in R \Rightarrow R$ is symmetric. Again,

(x, y)

$\in R$ and $(y, z) \in R$ then $(x, z) \in R \Rightarrow R$ is transitive. Therefore, R is reflexive, symmetric and transitive.

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

(c) $R = \{(x, y) : x \text{ is exactly 7 cm taller than } y\}$

x can not be taller than x , so R is not reflexive.

x is taller than y then y can not be taller than x , so R is not symmetric.

Again, x is 7 cm taller than y and y is 7 cm taller than z , then x can not be 7 cm taller than z , so R is not transitive.

Therefore, R is neither reflexive, nor symmetric and nor transitive.

(d) $R = \{(x, y) : x \text{ is wife of } y\}$

x is not wife of x , so R is not reflexive.

x is wife of y but y is not wife of x , so R is not symmetric.

Again, x is wife of y and y is wife of z then x can not be wife of z , so R is not transitive.

Therefore, R is neither reflexive, nor symmetric and nor transitive.

(e) $R = \{(x, y) : x \text{ is father of } y\}$

x is not father of x , so R is not reflexive.

x is father of y but y is not father of x , so R is not symmetric.

Again, x is father of y and y is father of z then x cannot be father of z , so R is not transitive.

Therefore, R is neither reflexive, nor symmetric and nor transitive.

2. Show that the relation R in the set R of real numbers, defined as $R = \{(a, b) : a \leq b^2\}$ is neither reflexive nor symmetric nor transitive.

Solution:

$R = \{(a, b) : a \leq b^2\}$, Relation R is defined as the set of real numbers.

(a, a)

$\in R$ then $a \leq a^2$, which is false. R is not reflexive. $(a, b) = (b, a)$

$\in R$ then $a \leq b^2$ and $b \leq a^2$, it is false statement. R is not symmetric. Now, $a \leq b^2$ and $b \leq c^2$, then a

$\leq c^2$, which is false. R is not transitive

Therefore, R is neither reflexive, nor symmetric and nor transitive.

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

3. Check whether the relation R defined in the set $\{1, 2, 3, 4, 5, 6\}$ as $R = \{(a, b) : b = a + 1\}$ is reflexive, symmetric or transitive.

Solution: $R = \{(a, b) : b = a + 1\}$

$R = \{(1, 2), (2, 3), (3, 4), (4, 5), (5, 6)\}$

When $b = a$, $a = a + 1$: which is false, So R is not reflexive.

If $(a, b) = (b, a)$, then $b = a + 1$ and $a = b + 1$: Which is false, so R is not symmetric.

Now, if (a, b) , (b, c) and (a, c) belongs to R then

$b = a + 1$ and $c = b + 1$ which implies $c = a + 2$: Which is false, so R is not transitive.

Therefore, R is neither reflexive, nor symmetric and nor transitive.

Q 4: Show that the relation R in R defined as $R = \{(a, b) : a \leq b\}$, is reflexive and transitive but not symmetric.

Solution:

Given relation is $R = \{(a, b) : a \leq b\}$

We know,

As $a \leq a$, so $(a, a) \in R$, therefore R is a reflexive relation.

As $a \leq b$ and $b \leq c$, then $a \leq c$, so $(a, b) \in R$, $(b, c) \in R$ and $(a, c) \in R$, therefore R is a transitive relation.

As $a \leq b$, then $a \geq b$ is not true,

For example, $(1, 2) \in R$ because $1 \leq 2$ is true but $(2, 1) \notin R$ because $2 \leq 1$ is false, therefore R is not symmetric relation.

Hence, R is reflexive and transitive but not symmetric.

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

Q 5: Check whether the relation R in R defined as $R = \{(a, b) : a \leq b^3\}$ is reflexive, symmetric or transitive.

Solution:

Given relation is $R = \{(a, b) : a \leq b^3\}$

For reflexive relation, $(a, a) \in R$ and $a \leq a^3$ but this is not always true.

Let $a = \frac{1}{2}$, $b = \frac{1}{2}$

$(\frac{1}{2}, \frac{1}{2}) \notin R$ as $\frac{1}{2} \leq (\frac{1}{2})^3$ is false.

Therefore R is not a reflexive relation.

For symmetric relation, if $(a, b) \in R$, then $(b, a) \in R$

Let $a = 2$, $b = 12$

$(2, 12) \in R$ as $2 \leq 12^3$ is true but $(12, 2) \notin R$ as $12 \leq 2^3$ is false.

Therefore R is not a symmetric relation.

For transitive relation, if $(a, b) \in R$, $(b, c) \in R$, then $(a, c) \in R$

Let $a = 12$, $b = 3$, $c = 2$

$(12, 3) \in R$ as $12 \leq 3^3$ is true, $(3, 2) \in R$ as $3 \leq 2^3$ is true but $(12, 2) \notin R$ as $12 \leq 2^3$ is false.

Therefore R is not a transitive relation.

Hence, R is neither reflexive, symmetric nor transitive.

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

6. Show that the relation R in the set {1, 2, 3} given by $R = \{(1, 2), (2, 1)\}$ is symmetric but neither reflexive nor transitive.

Solution:

$$R = \{(1, 2), (2, 1)\}$$

$$(x, x)$$

$\notin R$. R is not reflexive. $(1, 2)$

$\in R$ and $(2, 1) \in R$. R is symmetric. Again, (x, y)

$\in R$ and $(y, z) \in R$ then (x, z) does not imply to R. R is not transitive. Therefore, R is symmetric but neither reflexive nor transitive.

7. Show that the relation R in the set A of all the books in a library of a college, given by $R = \{(x, y) : x \text{ and } y \text{ have same number of pages}\}$ is an equivalence relation.

Solution:

Books x and x have same number of pages. (x, x)

$\in R$. R is reflexive. If (x, y)

$\in R$ and $(y, x) \in R$, so R is symmetric. Because, Books x and y have same number of pages and Books y and x have same number of pages.

Again, (x, y)

$\in R$ and $(y, z) \in R$ and $(x, z) \in R$. R is transitive. Therefore, R is an equivalence relation.

8. Show that the relation R in the set $A = \{1, 2, 3, 4, 5\}$ given by

$R = \{(a, b) : |a - b| \text{ is even}\}$, is an equivalence relation. Show that all the elements of $\{1, 3, 5\}$ are related to each other and all the elements of $\{2, 4\}$ are related to each other. But no element of $\{1, 3, 5\}$ is related to any element of $\{2, 4\}$.

Solution:

$$A = \{1, 2, 3, 4, 5\} \text{ and } R = \{(a, b) : |a - b| \text{ is even}\}$$

$$\text{We get, } R = \{(1, 3), (1, 5), (3, 5), (2, 4)\}$$

For (a, a) , $|a - b| = |a - a| = 0$ is even. Therefore, R is reflexive.

If $|a - b|$ is even, then $|b - a|$ is also even. R is symmetric.

Again, if $|a - b|$ and $|b - c|$ is even then $|a - c|$ is also even. R is transitive.

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

Therefore, R is an equivalence relation.

(b) We have to show that, Elements of $\{1, 3, 5\}$ are related to each other.

$$|1 - 3| = 2$$

$$|3 - 5| = 2$$

$$|1 - 5| = 4$$

All are even numbers.

Elements of $\{1, 3, 5\}$ are related to each other.

Similarly, $|2 - 4| = 2$ (even number), elements of $\{2, 4\}$ are related to each other.

Hence no element of $\{1, 3, 5\}$ is related to any element of $\{2, 4\}$.

9. Show that each of the relation R in the set $A = \{x$

$\in \mathbb{Z} : 0 \leq x \leq 12\}$, given by (i) $R = \{(a, b) : |a - b| \text{ is a multiple of } 4\}$ (ii) $R = \{(a, b) : a = b\}$ is an equivalence relation. Find the set of all elements related to 1 in each case.

Solution:

(i) $A = \{x$

$\in \mathbb{Z} : 0 \leq x \leq 12\}$ So, $A = \{0, 1, 2, 3, \dots, 12\}$

Now $R = \{(a, b) : |a - b| \text{ is a multiple of } 4\}$

$$R = \{(4, 0), (0, 4), (5, 1), (1, 5), (6, 2), (2, 6), \dots, (12, 9), (9, 12), \dots, (8, 0), (0, 8), \dots, (8, 4), (4, 8), \dots, (12, 12)\}$$

Here, $(x, x) = |4-4| = |8-8| = |12-12| = 0$: multiple of 4.

R is reflexive.

$|a - b|$ and $|b - a|$ are multiple of 4. (a, b)

$\in R$ and $(b, a) \in R$. R is symmetric.

And $|a - b|$ and $|b - c|$ then $|a - c|$ are multiple of 4. (a, b)

$\in R$ and $(b, c) \in R$ and $(a, c) \in R$ R is transitive.

Hence R is an equivalence relation.

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

(ii) Here, $(a, a) = a = a$.

(a, a)

$\in R$. So R is reflexive. $a = b$ and $b = a$. (a, b)

$\in R$ and $(b, a) \in R$. R is symmetric.

And $a = b$ and $b = c$ then $a = c$. (a, b)

$\in R$ and $(b, c) \in R$ and $(a, c) \in R$ R is transitive.

Hence R is an equivalence relation.

Now set of all elements related to 1 in each case is

(i) Required set = $\{1, 5, 9\}$

(ii) Required set = $\{1\}$

10. Give an example of a relation. Which is

(i) Symmetric but neither reflexive nor transitive.

(ii) Transitive but neither reflexive nor symmetric.

(iii) Reflexive and symmetric but not transitive.

(iv) Reflexive and transitive but not symmetric.

(v) Symmetric and transitive but not reflexive.

Solution:

(i) Consider a relation $R = \{(1, 2), (2, 1)\}$ in the set $\{1, 2, 3\}$

(x, x)

$\notin R$. R is not reflexive. $(1, 2)$

$\in R$ and $(2, 1) \in R$. R is symmetric. Again, (x, y)

$\in R$ and $(y, z) \in R$ then (x, z) does not imply to R . R is not transitive. Therefore, R is symmetric but neither

(ii) Relation $R = \{(a, b): a > b\}$

$a > a$ (false statement).

Also $a > b$ but $b > a$ (false statement) and

If $a > b$ but $b > c$, this implies $a > c$

Therefore, R is transitive, but neither reflexive nor symmetric.

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

(iii) $R = \{a, b\}$: a is friend of b}

a is friend of a. R is reflexive.

Also a is friend of b and b is friend of a. R is symmetric.

Also if a is friend of b and b is friend of c then a cannot be friend of c. R is not transitive.

Therefore, R is reflexive and symmetric but not transitive.

(iv) Say R is defined in R as $R = \{(a, b) : a \leq b\}$

$a \leq a$: which is true, (a, a)

$\in R$, So R is reflexive. $a \leq b$ but $b \leq a$ (false): (a, b)

$\in R$ but $(b, a) \notin R$, So R is not symmetric. Again, $a \leq b$ and $b \leq c$ then $a \leq c$: (a, b)

$\in R$ and (b, c) and $(a, c) \in R$, So R is transitive. Therefore, R is reflexive and transitive but not symmetric.

(v) $R = \{(a, b) : a \text{ is sister of } b\}$ (suppose a and b are female)

a is not sister of a. R is not reflexive.

a is sister of b and b is sister of a. R is symmetric.

Again, a is sister of b and b is sister of c then a is sister of c.

Therefore, R is symmetric and transitive but not reflexive.

11. Show that the relation R in the set A of points in a plane given by

$R = \{(P, Q) : \text{distance of the point P from the origin is same as the distance of the point Q from the origin}\}$, is an equivalence relation. Further, show that the set of all points related to a point $P \neq (0, 0)$ is the circle passing through P with origin as centre.

Solution: $R = \{(P, Q) : \text{distance of the point P from the origin is the same as the distance of the point Q from the origin}\}$

Say "O" is origin Point.

Since the distance of the point P from the origin is always the same as the distance of the same point P from the origin.

$OP = OP$

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

So $(P, P) \in R$. R is reflexive.

Distance of the point P from the origin is the same as the distance of the point Q from the origin

$$OP = OQ \text{ then } OQ = OP$$

R is symmetric.

Also $OP = OQ$ and $OQ = OR$ then $OP = OR$. R is transitive.

Therefore, R is an equivalent relation.

12. Show that the relation R defined in the set A of all triangles as $R = \{(T_1, T_2) : T_1 \text{ is similar to } T_2\}$, is equivalence relation. Consider three right angle triangles T_1 with sides 3, 4, 5, T_2 with sides 5, 12, 13 and T_3 with sides 6, 8, 10. Which triangles among T_1 , T_2 and T_3 are related?

Solution:

Case I:

T_1, T_2 are triangle.

$$R = \{(T_1, T_2) : T_1 \text{ is similar to } T_2\}$$

Check for reflexive:

As We know that each triangle is similar to itself, so (T_1, T_1)

$\in R$ R is reflexive.

Check for symmetric:

Also two triangles are similar, then T_1 is similar to T_2 and T_2 is similar to T_1 , so (T_1, T_2)

$$\in R \text{ and } (T_2, T_1)$$

$\in R$ R is symmetric.

Check for transitive:

Again, if then T_1 is similar to T_2 and T_2 is similar to T_3 , then T_1 is similar to T_3 , so (T_1, T_2)

$$\in R \text{ and } (T_2, T_3)$$

$\in R$ and $(T_1, T_3) \in R$ R is transitive

Therefore, R is an equivalent relation.

Case 2: It is given that T_1, T_2 and T_3 are right angled triangles.

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

T1 with sides 3, 4, 5

T2 with sides 5, 12, 13 and

T3 with sides 6, 8, 10

Since, two triangles are similar if corresponding sides are proportional.

Therefore, $3/6 = 4/8 = 5/10 = 1/2$

Therefore, T1 and T3 are related.

13. Show that the relation R defined in the set A of all polygons as $R = \{(P1, P2) : P1 \text{ and } P2 \text{ have same number of sides}\}$, is an equivalence relation. What is the set of all elements in A related to the right angle triangle T with sides 3, 4 and 5?

Solution:

Case I:

$R = \{(P1, P2) : P1 \text{ and } P2 \text{ have same number of sides}\}$

Check for reflexive:

P1 and P1 have same number of sides, So R is reflexive.

Check for symmetric:

P1 and P2 have same number of sides then P2 and P1 have same number of sides, so $(P1, P2)$

$\in R$ and $(P2, P1) \in R$ R is symmetric.

Check for transitive:

Again, P1 and P2 have same number of sides, and P2 and P3 have same number of sides, then also P1 and P3 have same number of sides .

So $(P1, P2)$

$\in R$ and $(P2, P3) \in R$ and $(P1, P3) \in R$ R is transitive

Therefore, R is an equivalent relation.

Since 3, 4, 5 are the sides of a triangle, the triangle is right angled triangle. Therefore, the set A is the set of right angled triangle.

14. Let L be the set of all lines in XY plane and R be the relation in L defined as $R = \{(L1, L2) : L1 \text{ is parallel to } L2\}$. Show that R is an equivalence relation. Find the set of all lines related to the line $y = 2x + 4$.

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

Solution:

L_1 is parallel to itself i.e., $(L_1, L_1) \in R$.

R is reflexive.

Now, let $(L_1, L_2) \in R$.

L_1 is parallel to L_2 and L_2 is parallel to L_1 .

$(L_2, L_1) \in R$, Therefore, R is symmetric.

Now, let $(L_1, L_2), (L_2, L_3) \in R$.

L_1 is parallel to L_2 . Also, L_2 is parallel to L_3 .

L_1 is parallel to L_3 .

Therefore, R is transitive.

Hence, R is an equivalence relation.

Again, The set of all lines related to the line $y = 2x + 4$, is the set of all its parallel lines.

Slope of given line is $m = 2$.

As we know slope of all parallel lines are same.

Hence, the set of all related to $y = 2x + 4$ is $y = 2x + k$, where k

$\in \mathbb{R}$. **15. Let R be the relation in the set $\{1, 2, 3, 4\}$ given by $R = \{(1, 2), (2, 2), (1, 1), (4, 4), (1, 3), (3, 3), (3, 2)\}$. Choose the correct answer.**

(A) R is reflexive and symmetric but not transitive.

(B) R is reflexive and transitive but not symmetric.

(C) R is symmetric and transitive but not reflexive.

(D) R is an equivalence relation.

Solution:

Let R be the relation in the set $\{1, 2, 3, 4\}$ given by $R = \{(1, 2), (2, 2), (1, 1), (4, 4), (1, 3), (3, 3), (3, 2)\}$.

Step 1: $(1, 1), (2, 2), (3, 3), (4, 4)$

$\in R$. R is reflexive. Step 2: $(1, 2)$

$\in R$ but $(2, 1) \notin R$. R is not symmetric. Step 3: Consider any set of points, $(1, 3)$

$\in R$ and $(3, 2) \in R$ then $(1, 2) \in R$. So R is transitive.

Option (B) is correct.

16. Let R be the relation in the set \mathbb{N} given by $R = \{(a, b) : a = b - 2, b > 6\}$. Choose the correct answer.

(A) $(2, 4)$

$\in R$ **(B) $(3, 8) \in R$ (C) $(6, 8) \in R$ (D) $(8, 7) \in R$**

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

Solution: $R = \{(a, b) : a = b - 2, b > 6\}$

(A) Incorrect : Value of $b = 4$, not true.

(B) Incorrect : $a = 3$ and $b = 8 > 6$

$a = b - 2 \Rightarrow 3 = 8 - 2$ and $3 = 6$, which is false.

(C) Correct: $a = 6$ and $b = 8 > 6$

$a = b - 2 \Rightarrow 6 = 8 - 2$ and $6 = 6$, which is true.

(D) Incorrect : $a = 8$ and $b = 7 > 6$

$a = b - 2 \Rightarrow 8 = 7 - 2$ and $8 = 5$, which is false.

Therefore, option (C) is correct.

R but (2, 1)

$\notin R$

\in

Exercise 1.2

1. Show that the function $f : \mathbb{R} \setminus \{0\} \rightarrow \mathbb{R} \setminus \{0\}$ defined by $f(x) = 1/x$ is one-one and onto, where $\mathbb{R} \setminus \{0\}$ is the set of all non-zero real numbers. Is the function one-one and onto if $\mathbb{R} \setminus \{0\}$ is replaced by \mathbb{N} with co-domain being same as $\mathbb{R} \setminus \{0\}$?

Solution:

Given: $f : \mathbb{R} \setminus \{0\} \rightarrow \mathbb{R} \setminus \{0\}$ defined by $f(x) = 1/x$

Check for One-One

$$\text{If } f(x_1) = f(x_2) \text{ then } \frac{1}{x_1} = \frac{1}{x_2}$$

$$\text{This implies } x_1 = x_2$$

Therefore, f is one-one function.

Check for onto

$$f(x) = 1/x$$

$$\text{or } y = 1/x$$

$$\text{or } x = 1/y$$

$$f(1/y) = y$$

Therefore, f is onto function.

Again, If $f(x_1) = f(x_2)$

Say, n_1, n_2

$$\in \mathbb{N}$$

$$\frac{1}{n_1} = \frac{1}{n_2}$$

$$= n_1 n_2$$

So $n_1 = n_2$

Therefore, f is one-one

Every real number belonging to co-domain may not have a pre-image in \mathbb{N} . for example, $1/3$ and $3/2$ are not belongs \mathbb{N} . So \mathbb{N} is not onto.

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

2. Check the injectivity and surjectivity of the following functions:

(i) $f : \mathbf{N} \rightarrow \mathbf{N}$ given by $f(x) = x^2$

(ii) $f : \mathbf{Z} \rightarrow \mathbf{Z}$ given by $f(x) = x^2$

(iii) $f : \mathbf{R} \rightarrow \mathbf{R}$ given by $f(x) = x^2$

(iv) $f : \mathbf{N} \rightarrow \mathbf{N}$ given by $f(x) = x^3$

(v) $f : \mathbf{Z} \rightarrow \mathbf{Z}$ given by $f(x) = x^3$

Solution:

(i) $f : \mathbf{N} \rightarrow \mathbf{N}$ given by $f(x) = x^2$

For x, y

$\in \mathbf{N} \Rightarrow f(x) = f(y)$ which implies $x^2 = y^2 \quad x = y$ Therefore f is injective.

There are such numbers of co-domain which have no image in domain \mathbf{N} .

Say, 3

$\in \mathbf{N}$, but there is no pre-image in domain of f . such that $f(x) = x^2 = 3$. f is not surjective.

Therefore, f is injective but not surjective.

(ii) Given, $f : \mathbf{Z} \rightarrow \mathbf{Z}$ given by $f(x) = x^2$

Here, $\mathbf{Z} = \{0, \pm 1, \pm 2, \pm 3, \pm 4, \dots\}$

$f(-1) = f(1) = 1$

But -1 not equal to 1 .

f is not injective.

There are many numbers of co-domain which have no image in domain \mathbf{Z} .

For example, -3

\in co-domain \mathbf{Z} , but $-3 \notin$ domain \mathbf{Z} f is not surjective.

Therefore, f is neither injective nor surjective.

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

(iii) $f : \mathbf{R} \rightarrow \mathbf{R}$ given by $f(x) = x^2$

$$f(-1) = f(1) = 1$$

But -1 not equal to 1.

f is not injective.

There are many numbers of co-domain which have no image in domain \mathbf{R} .

For example, -3

\in co-domain \mathbf{R} , but there does not exist any x in domain \mathbf{R} where $x^2 = -3$ f is not surjective.

Therefore, f is neither injective nor surjective.

(iv) $f : \mathbf{N} \rightarrow \mathbf{N}$ given by $f(x) = x^3$

For x, y

$$\in \mathbf{N} \Rightarrow f(x) = f(y) \text{ which implies } x^3 = y^3 \quad x = y \text{ Therefore } f \text{ is injective.}$$

There are many numbers of co-domain which have no image in domain \mathbf{N} .

For example, 4

\in co-domain \mathbf{N} , but there does not exist any x in domain \mathbf{N} where $x^3 = 4$. f is not surjective.

Therefore, f is injective but not surjective.

(v) $f : \mathbf{Z} \rightarrow \mathbf{Z}$ given by $f(x) = x^3$

For x, y

$$\in \mathbf{Z} \Rightarrow f(x) = f(y) \text{ which implies } x^3 = y^3 \quad x = y \text{ Therefore } f \text{ is injective.}$$

There are many numbers of co-domain which have no image in domain \mathbf{Z} .

For example, 4

\in co-domain \mathbf{N} , but there does not exist any x in domain \mathbf{Z} where $x^3 = 4$. f is not surjective.

Therefore, f is injective but not surjective.

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

3. Prove that the Greatest Integer Function $f : \mathbb{R} \rightarrow \mathbb{R}$, given by $f(x) = [x]$, is neither one-one nor onto, where $[x]$ denotes the greatest integer less than or equal to x .

Solution:

Function $f : \mathbb{R} \rightarrow \mathbb{R}$, given by $f(x) = [x]$
 $f(x) = 1$, because $1 \leq x \leq 2$

$$f(1.2) = [1.2] = 1$$

$$f(1.9) = [1.9] = 1$$

But $1.2 \neq 1.9$

f is not one-one.

There is no fraction proper or improper belonging to co-domain of f has any pre-image in its domain.

For example, $f(x) = [x]$ is always an integer

for 0.7 belongs to \mathbb{R} there does not exist any x in domain \mathbb{R} where $f(x) = 0.7$ f is not onto.

Hence proved, the Greatest Integer Function is neither one-one nor onto.

4. Show that the Modulus Function $f : \mathbb{R} \rightarrow \mathbb{R}$, given by $f(x) = |x|$, is neither one-one nor onto, where $|x|$ is x , if x is positive or 0 and $|x|$ is $-x$, if x is negative.

Solution:

$f : \mathbb{R} \rightarrow \mathbb{R}$, given by $f(x) = |x|$, defined as

$$f(x) = |x| = \begin{cases} x, & \text{if } x \geq 0 \\ -x, & \text{if } x < 0 \end{cases}$$

f contains values like $(-1, 1), (1, 1), (-2, 2), (2, 2)$

$$f(-1) = f(1), \text{ but } -1 \neq 1$$

f is not one-one.

\mathbb{R} contains some negative numbers which are not images of any real number since $f(x) = |x|$ is always non-negative. So f is not onto.

Hence, Modulus Function is neither one-one nor onto.

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

5. Show that the Signum Function $f : \mathbb{R} \rightarrow \mathbb{R}$, given by

$$f(x) = \begin{cases} 1, & \text{if } x > 0 \\ 0, & \text{if } x = 0 \\ -1, & \text{if } x < 0 \end{cases}$$

is neither one-one nor onto.

Solution: Signum Function $f : \mathbb{R} \rightarrow \mathbb{R}$, given by

$$f(x) = \begin{cases} 1, & \text{if } x > 0 \\ 0, & \text{if } x = 0 \\ -1, & \text{if } x < 0 \end{cases}$$

$$f(1) = f(2) = 1$$

This implies, for $n > 0, f$

$$f(x_1) = f(x_2) = 1 \quad x_1 \neq x_2$$

f is not one-one.

$f(x)$ has only 3 values, $(-1, 0, 1)$. Other than these 3 values of co-domain \mathbb{R} has no any pre-image its domain.

f is not onto.

Hence, Signum Function is neither one-one nor onto.

6. Let $A = \{1, 2, 3\}$, $B = \{4, 5, 6, 7\}$ and let $f = \{(1, 4), (2, 5), (3, 6)\}$ be a function from A to B . Show that f is one-one.

Solution:

$$A = \{1, 2, 3\}$$

$$B = \{4, 5, 6, 7\} \text{ and}$$

$$f = \{(1, 4), (2, 5), (3, 6)\}$$

$$f(1) = 4, f(2) = 5 \text{ and } f(3) = 6$$

Here, also distinct elements of A have distinct images in B .

Therefore, f is one-one.

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

7. In each of the following cases, state whether the function is one-one, onto or bijective. Justify your answer.

(i) $f : \mathbb{R} \rightarrow \mathbb{R}$ defined by $f(x) = 3 - 4x$

(ii) $f : \mathbb{R} \rightarrow \mathbb{R}$ defined by $f(x) = 1 + x^2$

Solution:

(i) $f : \mathbb{R} \rightarrow \mathbb{R}$ defined by $f(x) = 3 - 4x$

If x_1, x_2

$\in \mathbb{R}$ then $f(x_1) = 3 - 4x_1$ and

$f(x_2) = 3 - 4x_2$

If $f(x_1) = f(x_2)$ then $x_1 = x_2$

Therefore, f is one-one.

Again,

$f(x) = 3 - 4x$

or $y = 3 - 4x$

or $x = (3-y)/4 \in \mathbb{R}$

$f((3-y)/4) = 3 - 4((3-y)/4) = y$

f is onto.

Hence f is onto or bijective.

(ii) $f : \mathbb{R} \rightarrow \mathbb{R}$ defined by $f(x) = 1 + x^2$

If x_1, x_2

$\in \mathbb{R}$ then $f(x) = 1 + x$

x_1

and $f(x) = 1 + x$

x_2

If $f(x_1) = f(x_2)$ then x_1

x_2

$= x_2$ This implies $x_1 \neq x_2$

Therefore, f is not one-one

Again, if every element of co-domain is image of some element of Domain under f , such that

$f(x) = y$

$f(x) = 1 + x^2$

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

$$y = f(x) = 1 + x^2 \text{ _____}$$

$$\text{or } x = \text{ _____}$$

$$\pm \sqrt{1 - y}$$

Therefore, f(

$\sqrt{1-y}$) = $2 - y \neq y$ Therefore, f is not onto or bijective.

8. Let A and B be sets. Show that $f : A \times B \rightarrow B \times A$ such that $f(a, b) = (b, a)$ is bijective function.

Solution:

Step 1: Check for Injectivity:

Let (a_1, b_1) and (a_2, b_2)

$\in A \times B$ such that $f(a_1, b_1) = (a_2, b_2)$

This implies, (b_1, a_1) and (b_2, a_2)

$$b_1 = b_2 \text{ and } a_1 = a_2$$

$(a_1, b_1) = (a_2, b_2)$ for all (a_1, b_1) and (a_2, b_2)

$\in A \times B$ Therefore, f is injective.

Step 2: Check for Surjectivity:

Let (b, a) be any element of $B \times A$. Then a

$\in A$ and $b \in B$ This implies (a, b)

$\in A \times B$ For all (b, a)

$\in B \times A$, there exists $(a, b) \in A \times B$ Therefore, $f: A \times B \rightarrow B \times A$ is bijective function.

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

9. Let $f : \mathbb{N} \rightarrow \mathbb{N}$ be defined by

$$f(n) = \begin{cases} \frac{n+1}{2}, & \text{if } n \text{ is odd} \\ \frac{n}{2}, & \text{if } n \text{ is even} \end{cases} \quad \text{for all } n \in \mathbb{N}$$

State whether the function f is bijective. Justify your answer

Solution:

$$f(n) = \begin{cases} \frac{n+1}{2}, & \text{if } n \text{ is odd} \\ \frac{n}{2}, & \text{if } n \text{ is even} \end{cases} \quad \text{for all } n \in \mathbb{N}$$

For $n = 1, 2$

$$f(1) = (n+1)/2 = (1+1)/2 = 1 \text{ and}$$

$$f(2) = (n)/2 = (2)/2 = 1$$

$$f(1) = f(2), \text{ but } 1 \neq 2$$

f is not one-one.

For a natural number, “ a ” in co-domain \mathbb{N}

If n is odd

$$n = 2k + 1 \text{ for } k$$

$$\in \mathbb{N}, \text{ then } 4k + 1 \in \mathbb{N} \text{ such that } f(4k+1) = (4k+1+1)/2 = 2k + 1$$

If n is even

$$n = 2k \text{ for some } k$$

$$\in \mathbb{N} \text{ such that } f(4k) = 4k/2 = 2k \text{ } f \text{ is onto}$$

Therefore, f is onto but not bijective function.

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

10. Let $A = \mathbb{R} - \{3\}$ and $B = \mathbb{R} - \{1\}$. Consider the function $f : A \rightarrow B$ defined by

$$f(x) = \frac{(x-2)}{(x-3)}$$

Is f one-one and onto? Justify your answer.

Solution: $A = \mathbb{R} - \{3\}$ and $B = \mathbb{R} - \{1\}$

$f : A \rightarrow B$ defined by $f(x) = \frac{(x-2)}{(x-3)}$

Let $(x, y) \in A$ then

$$f(x) = \frac{x-2}{x-3} \quad \text{and} \quad f(y) = \frac{y-2}{y-3}$$

$$\frac{x-2}{x-3} = \frac{y-2}{y-3}$$

$$(x-2)(y-3) = (y-2)(x-3)$$

$$xy - 3x - 2y + 6 = xy - 3y - 2x + 6$$

$$-3x - 2y = -3y - 2x$$

$$-3x + 2x = -3y + 2y$$

$$-x = -y$$

$$x = y$$

Again, $f(x) = \frac{(x-2)}{(x-3)}$

or $y = f(x) = \frac{(x-2)}{(x-3)}$

$$y = \frac{(x-2)}{(x-3)}$$

$$y(x-3) = x-2$$

$$xy - 3y = x - 2$$

$$x(y-1) = 3y - 2$$

$$\text{or } x = \frac{(3y-2)}{(y-1)}$$

$$3y - 2$$

$$-1 - 2y \text{ Now, } f\left(\frac{(3y-2)}{(y-1)}\right) = \frac{3y-2}{y-1} = y$$

$$-1 - 3y$$

$$f(x) = y$$

Therefore, f is onto function.

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

11. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be defined as $f(x) = x^4$. Choose the correct answer.

- (A) f is one-one onto (B) f is many-one onto
(C) f is one-one but not onto (D) f is neither one-one nor onto.**

Solution:

$f : \mathbb{R} \rightarrow \mathbb{R}$ be defined as $f(x) = x^4$

let x and y belongs to \mathbb{R} such that, $f(x) = f(y)$

$$x^4 = y^4 \text{ or } x = \pm y$$

f is not one-one function.

$$\text{Now, } y = f(x) = x^4 \text{ Or } x = \pm y^{1/4}$$

$$f(y^{1/4}) = y \text{ and } f(-y^{1/4}) = -y$$

Therefore, f is not onto function.

Option D is correct.

12. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be defined as $f(x) = 3x$. Choose the correct answer.

- (A) f is one-one onto (B) f is many-one onto
(C) f is one-one but not onto (D) f is neither one-one nor onto.**

Solution: $f : \mathbb{R} \rightarrow \mathbb{R}$ be defined as $f(x) = 3x$

let x and y belongs to \mathbb{R} such that $f(x) = f(y)$

$$3x = 3y \text{ or } x = y$$

f is one-one function.

$$\text{Now, } y = f(x) = 3x$$

$$\text{Or } x = y/3$$

$$f(x) = f(y/3) = y$$

Therefore, f is onto function.

Option (A) is correct.

Exercise 1.3

Page No: 18

1. Let $f : \{1, 3, 4\} \rightarrow \{1, 2, 5\}$ and $g : \{1, 2, 5\} \rightarrow \{1, 3\}$ be given by $f = \{(1, 2), (3, 5), (4, 1)\}$ and $g = \{(1, 3), (2, 3), (5, 1)\}$. Write down gof .

Solution:

Given function, $f : \{1, 3, 4\} \rightarrow \{1, 2, 5\}$ and $g : \{1, 2, 5\} \rightarrow \{1, 3\}$ be given by

$f = \{(1, 2), (3, 5), (4, 1)\}$ and $g = \{(1, 3), (2, 3), (5, 1)\}$

Find gof .

At $f(1) = 2$ and $g(2) = 3$, gof is

$$gof(1) = g(f(1)) = g(2) = 3$$

At $f(3) = 5$ and $g(5) = 1$, gof is

$$gof(3) = g(f(3)) = g(5) = 1$$

At $f(4) = 1$ and $g(1) = 3$, gof is

$$gof(4) = g(f(4)) = g(1) = 3$$

Therefore, $gof = \{(1,3), (3,1), (4,3)\}$

2. Let f, g and h be functions from R to R . Show that

$$(f + g) oh = foh + goh$$

$$(f \cdot g) oh = (foh) \cdot (goh)$$

Solution:

$$\text{LHS} = (f + g) oh$$

$$= (f+g)(h(x))$$

$$= f(h(x)) + g(h(x))$$

$$= foh + goh$$

$$= \text{RHS}$$

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

Again,

$$\text{LHS} = (f \circ g) \circ h$$

$$= f \circ (g \circ h)$$

$$= f(h(x)) \circ g(h(x))$$

$$= (f \circ h) \circ (g \circ h)$$

$$= \text{RHS}$$

3. Find $g \circ f$ and $f \circ g$, if

(i) $f(x) = |x|$ and $g(x) = |5x - 2|$

(ii) $f(x) = 8x^3$ and $g(x) = x^{1/3}$.

Solution:

(i) $f(x) = |x|$ and $g(x) = |5x - 2|$

$$g \circ f = (g \circ f)(x) = g(f(x)) = g(|x|) = |5|x| - 2|$$

$$f \circ g = (f \circ g)(x) = f(g(x)) = f(|5x - 2|) = ||5x - 2|| = |5x - 2|$$

(ii) $f(x) = 8x^3$ and $g(x) = x^{1/3}$.

$$g \circ f = (g \circ f)(x) = g(f(x)) = g(8x^3) = (8x^3)^{1/3} = 2x$$

$$f \circ g = (f \circ g)(x) = f(g(x)) = f(x^{1/3}) = 8(x^{1/3})^3 = 8x$$

(4x+3)⁴ 4. If $f(x) = \frac{1}{x-2/3}$, Show that $f \circ f(x) = x$, for all $x \neq 2/3$. What is the inverse of f .

(6x-4)

Solution:

$$(4x+3)f(x) = \frac{1}{x-2/3},$$

$$(6x-4)$$

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

$$\begin{aligned} &= \frac{4\left(\frac{4x+3}{6x-4}\right)+3}{6\left(\frac{4x+3}{6x-4}\right)-4} \\ &= \frac{16x+12+18x-12}{24x+18-24x+16} \\ &= \frac{34x}{34} \\ &= x \end{aligned}$$

Therefore, $f \circ f(x) = x$ for all $x \neq 2/3$.

Again, $f \circ f = I$

The inverse of the given function, f is f .

5. State with reason whether following functions have inverse

(i) $f : \{1, 2, 3, 4\} \rightarrow \{10\}$ with
 $f = \{(1, 10), (2, 10), (3, 10), (4, 10)\}$

(ii) $g : \{5, 6, 7, 8\} \rightarrow \{1, 2, 3, 4\}$ with
 $g = \{(5, 4), (6, 3), (7, 4), (8, 2)\}$

(iii) $h : \{2, 3, 4, 5\} \rightarrow \{7, 9, 11, 13\}$ with
 $h = \{(2, 7), (3, 9), (4, 11), (5, 13)\}$

Solution:

(i) $f : \{1, 2, 3, 4\} \rightarrow \{10\}$ with $f = \{(1, 10), (2, 10), (3, 10), (4, 10)\}$

f has many-one function like $f(1) = f(2) = f(3) = f(4) = 10$, therefore f has no inverse.

(ii) $g : \{5, 6, 7, 8\} \rightarrow \{1, 2, 3, 4\}$ with $g = \{(5, 4), (6, 3), (7, 4), (8, 2)\}$

g has many-one function like $g(5) = g(7) = 4$, therefore g has no inverse.

(iii) $h : \{2, 3, 4, 5\} \rightarrow \{7, 9, 11, 13\}$ with $h = \{(2, 7), (3, 9), (4, 11), (5, 13)\}$

All elements have different images under h . So h is one-one onto function, therefore, h has an inverse.

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

6. Show that $f : [-1, 1] \rightarrow \mathbb{R}$, given by $f(x) = x/(x+2)$ is one-one. Find the inverse of the function $f : [-1, 1] \rightarrow \text{Range } f$.

(Hint: For y

$\in \text{Range } f$, $y = f(x) = x/(x+2)$, for some x in $[-1, 1]$, i.e., $x = 2y/(1-y)$. Solution:

Given function: $f(x) = x/(x+2)$

Let x, y

$\in [-1, 1]$ Let $f(x) = f(y)$

$$x/(x+2) = y/(y+2)$$

$$xy + 2x = xy + 2y$$

$$x = y$$

f is one-one.

Again,

Since $f : [-1, 1] \rightarrow \text{Range } f$ is onto

$$\text{say, } y = x/(x+2)$$

$$yx + 2y = x$$

$$x(1 - y) = 2y$$

$$\text{or } x = 2y/(1-y)$$

$$x = f^{-1}(y) = 2y/(1-y); y \text{ not equal to } 1$$

f is onto function, and $f^{-1}(x) = 2x/(1-x)$.

7. Consider $f : \mathbb{R} \rightarrow \mathbb{R}$ given by $f(x) = 4x + 3$. Show that f is invertible. Find the inverse of f .

Solution:

Consider $f : \mathbb{R} \rightarrow \mathbb{R}$ given by $f(x) = 4x + 3$

Say, x, y

$\in \mathbb{R}$ Let $f(x) = f(y)$ then

$$4x + 3 = 4y + 3$$

$$x = y$$

f is one-one function.

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

Let y

\in Range of f $y = 4x + 3$

or $x = (y-3)/4$

Here, $f((y-3)/4) = 4((y-3)/4) + 3 = y$

This implies $f(x) = y$

So f is onto

Therefore, f is invertible.

Inverse of f is $x = f^{-1}(y) = (y-3)/4$.

8. Consider $f : \mathbb{R}^+ \rightarrow [4, \infty)$ given by $f(x) = x + 4$. Show that f is invertible with the inverse f^{-1} of f given by $f^{-1}(y) =$ _____

$\sqrt{y-4}$, where \mathbb{R}^+ is the set of all non-negative real numbers. Solution:

Consider $f : \mathbb{R}^+ \rightarrow [4, \infty)$ given by $f(x) = x + 4$

Let x, y

$\in \mathbb{R} \rightarrow [4, \infty)$ then

$f(x) = x + 4$ and

$f(y) = y + 4$

if $f(x) = f(y)$ then $x + 4 = y + 4$

or $x = y$

f is one-one.

Now $y = f(x) = x + 4$ or $x =$ _____
 $\sqrt{y-4}$ as $x > 0$

$f($ _____ _____

$\sqrt{y-4}) = (\sqrt{y-4})^2 + 4 = y$ $f(x) = y$

f is onto function.

Therefore, f is invertible and Inverse of f is $f^{-1}(y) =$ _____
 $\sqrt{y-4}$.

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

9. Consider $f : \mathbb{R}^+ \rightarrow [-5, \infty)$ given by $f(x) = 9x^2 + 6x - 5$. Show that f is invertible with

Solution: $f^{-1}(y) = \left(\frac{(\sqrt{y+6})-1}{3} \right)$

Consider $f : \mathbb{R}^+ \rightarrow [-5, \infty)$ given by $f(x) = 9x^2 + 6x - 5$

Consider $f : \mathbb{R}^+ \rightarrow [4, \infty)$ given by $f(x) = x + 4$

Let x, y

$\in \mathbb{R}^+ \rightarrow [-5, \infty)$ then

$$f(x) = 9x^2 + 6x - 5 \text{ and}$$

$$f(y) = 9y^2 + 6y - 5$$

$$\text{if } f(x) = f(y) \text{ then } 9x^2 + 6x - 5 = 9y^2 + 6y - 5$$

$$9(x^2 - y^2) + 6(x - y) = 0$$

$$9\{(x-y)(x+y)\} + 6(x - y) = 0$$

$$(x - y)(9(x + y) + 6) = 0$$

either $x - y = 0$ or $9(x + y) + 6 = 0$

Say $x - y = 0$, then $x = y$. So f is one-one.

$$\text{Now, } y = f(x) = 9x^2 + 6x - 5$$

Solving this quadratic equation, we have

$$x = \frac{-6 \pm \sqrt{36 + 36(y+5)}}{18}$$

$$\text{So, } f(x) = f\left(\frac{-6 + \sqrt{36 + 36(y+5)}}{18}\right) = 9\left(\frac{-6 + \sqrt{36 + 36(y+5)}}{18}\right)^2 + 6\left(\frac{-6 + \sqrt{36 + 36(y+5)}}{18}\right) - 5$$

$$= y + 7 - 2 - 5 = y$$

$\sqrt{y+6} + 2$ $y = f(x)$, therefore, f is onto.

$$f(x) \text{ is invertible and } f^{-1}(x) = \frac{\sqrt{y+6}-1}{3}$$

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

10. Let $f : X \rightarrow Y$ be an invertible function. Show that f has unique inverse.

(Hint: suppose g_1 and g_2 are two inverses of f . Then for all $y \in Y, f \circ g_1(y) = 1_Y(y) = f \circ g_2(y)$. Use one-one ness of f)

Solution:

Given, $f : X \rightarrow Y$ be an invertible function. And g_1 and g_2 are two inverses of f .

For all y

$\in Y$, we get

$$f \circ g_1(y) = 1_Y(y) = f \circ g_2(y)$$

$$f(g_1(y)) = f(g_2(y))$$

$$g_1(y) = g_2(y)$$

$$g_1 = g_2$$

Hence f has unique inverse.

11. Consider $f : \{1, 2, 3\} \rightarrow \{a, b, c\}$ given by $f(1) = a$, $f(2) = b$ and $f(3) = c$. Find f^{-1} and show that $(f^{-1})^{-1} = f$.

Solution:

Consider $f : \{1, 2, 3\} \rightarrow \{a, b, c\}$ given by $f(1) = a$, $f(2) = b$ and $f(3) = c$

$$\text{So } f = \{(a, 1), (b, 2), (c, 3)\}$$

$$\text{Hence } f^{-1}(a) = 1, f^{-1}(b) = 2 \text{ and } f^{-1}(c) = 3$$

$$\text{Now, } f^{-1} = \{(a, 1), (b, 2), (c, 3)\}$$

$$\text{Therefore, inverse of } f^{-1} = (f^{-1})^{-1} = \{(1, a), (2, b), (3, c)\} = f$$

$$\text{Hence } (f^{-1})^{-1} = f.$$

13. If $f: \mathbb{R} \rightarrow \mathbb{R}$ be given by $f(x) =$

$(3-x^3)^3$, then $f \circ f(x)$ is (A) $x^{1/3}$ (B) x^3 (C) x (D) $(3 - x^3)$

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

Solution:

$f: \mathbb{R} \rightarrow \mathbb{R}$ be given by $f(x) = \left(3 - x^3\right)^{\frac{1}{3}}$, then

$$\begin{aligned} f \circ f(x) &= f(f(x)) \\ &= f\left(\left(3 - x^3\right)^{\frac{1}{3}}\right) \\ &= \left[3 - \left(\left(3 - x^3\right)^{\frac{1}{3}}\right)^3\right]^{\frac{1}{3}} \\ &= \left[3 - (3 - x^3)\right]^{\frac{1}{3}} \\ &= \left(x^3\right)^{\frac{1}{3}} = x \end{aligned}$$

Option (C) is correct.

4x

14. Let $f: \mathbb{R} - \{-4/3\} \rightarrow \mathbb{R}$ be a function defined as $f(x) = \frac{4x}{3x+4}$. The inverse of f is the

$+4$ map $g: \text{Range } f \rightarrow \mathbb{R} - \{-4/3\}$ given by

(A) $g(y) = 3y/(3-4y)$ (B) $g(y) = 4y/(4-3y)$

(C) $g(y) = 4y/(3-4y)$ (D) $g(y) = 3y/(4-3y)$

Solution:

Let $f: \mathbb{R} - \{-4/3\} \rightarrow \mathbb{R}$ be a function defined as $f(x) = \frac{4x}{3x+4}$. And $\text{Range } f \rightarrow \mathbb{R} - \{-4/3\}$

$3x + 4$

$4xy = f(x) = \frac{4x}{3x+4}$

$3x + 4$

$y(3x + 4) = 4x$

$3xy + 4y = 4x$

$x(3y - 4) = -4y$

$x = 4y/(4-3y)$

Therefore, $f^{-1}(y) = g(y) = 4y/(4-3y)$. Option (B) is the correct answer.

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

1. Determine whether or not each of the definition of

*** given below gives a binary operation. In the event that**

*** is not a binary operation, give justification for this. (i) On Z^+ , define**

*** by $a * b = a - b$ (ii) On Z^+ , define**

*** by $a * b = ab$ (iii) On R , define**

*** by $a * b = ab^2$ (iv) On Z^+ , define**

*** by $a * b = |a - b|$ (v) On Z^+ , define**

*** by $a * b = a$ Solution:**

(i) On Z^+ , define

*** by $a * b = a - b$ On $Z^+ = \{1, 2, 3, 4, 5, \dots\}$**

Let $a = 1$ and $b = 2$

Therefore, a

*** $b = a - b = 1 - 2 = -1 \notin Z^+$ operation * is not a binary operation on Z^+ .**

(ii) On Z^+ , define

*** by $a * b = ab$ On $Z^+ = \{1, 2, 3, 4, 5, \dots\}$**

Let $a = 2$ and $b = 3$

Therefore, a

*** $b = a * b = 2 * 3 = 6 \in Z^+$ operation * is a binary operation on Z^+**

(iii) On R , define

*** by $a * b = ab^2$ $R = \{-\infty, \dots, -1, 0, 1, 2, \dots, \infty\}$**

Let $a = 1.2$ and $b = 2$

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

Therefore, a

* $b = ab^2 = (1.2) \times 2^2 = 4.8 \in \mathbb{R}$ Operation * is a binary operation on \mathbb{R} .

(iv) On \mathbb{Z}^+ , define

* **by $a * b = |a - b|$ On $\mathbb{Z}^+ = \{1, 2, 3, 4, 5, \dots\}$**

Let $a = 2$ and $b = 3$

Therefore, a

* $b = a * b = 2 * 3 = 6 \in \mathbb{Z}^+$ operation * is a binary operation on \mathbb{Z}^+

(v) On \mathbb{Z}^+ , define

* **by $a * b = a$ On $\mathbb{Z}^+ = \{1, 2, 3, 4, 5, \dots\}$**

Let $a = 2$ and $b = 1$

Therefore, a

* $b = 2 \in \mathbb{Z}^+$ Operation * is a binary operation on \mathbb{Z}^+ .

2. For each operation

* **defined below, determine whether * or associative. is binary, commutative**

(i) On \mathbb{Z} , define a

* **$b = a - b$ (ii) On \mathbb{Q} , define a**

* **$b = ab + 1$ (iii) On \mathbb{Q} , define a**

* **$b = ab/2$ (iv) On \mathbb{Z}^+ , define a**

* **$b = 2ab$ (v) On \mathbb{Z}^+ , define a**

* **$b = ab$ (vi) On $\mathbb{R} - \{-1\}$, define a**

* **$b = a/(b+1)$**

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

Solution:

(i) On \mathbb{Z} , define a

* **$b = a - b$** Step 1: Check for commutative

Consider

* is commutative, then a

* $b = b * a$ Which means, $a - b = b - a$ (not true)

Therefore,

* is not commutative. Step 2: Check for Associative.

Consider

* is associative, then (a

* $b) * c = a * (b * c)$ LHS = (a

* $b) * c = (a - b) * c = a - b - c$

RHS = $a * (b * c) = a - (b - c)$

= $a - (b - c)$

= $a - b + c$

This implies LHS \neq RHS

Therefore,

* is not associative. **(ii) On \mathbb{Q} , define a**

* **$b = ab + 1$** Step 1: Check for commutative

Consider

* is commutative, then a

* $b = b * a$ Which means, $ab + 1 = ba + 1$

or $ab + 1 = ab + 1$ (which is true)

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

a

* $b = b * a$ for all $a, b \in Q$ Therefore,

* is commutative. Step 2: Check for Associative.

Consider

* is associative, then (a

* $b) * c = a * (b * c)$ LHS = (a

* $b) * c = (ab + 1) * c = (ab + 1)c + 1$

= $abc + c + 1$

RHS = $a * (b * c) = a * (bc + 1)$

= $a(bc + 1) + 1$

= $abc + a + 1$

This implies $LHS \neq RHS$

Therefore,

* is not associative. **(iii) On Q, define a**

* **$b = ab/2$** Step 1: Check for commutative

Consider

* is commutative, then a

* $b = b * a$ Which means, $ab/2 = ba/2$

or $ab/2 = ab/2$ (which is true)

a

* $b = b * a$ for all $a, b \in Q$ Therefore,

* is commutative.

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

Step 2: Check for Associative.

Consider

* is associative, then (a

$$* b)^* c = a * (b * c) \text{ LHS} = (a$$

$$* b) * c = (ab/2) * c \text{ ab}$$

$$\frac{2 \times c}{2} =$$

$$= abc/4$$

$$\text{RHS} = a * (b * c) = a * (bc/2)$$

$$\times bca = \frac{2}{2}$$

$$= abc/4$$

This implies LHS = RHS

Therefore,

* is associative binary operation. **(iv) On Z^+ , define a**

* **$b = 2ab$** Step 1: Check for commutative

Consider

* is commutative, then a

$$* b = b * a \text{ Which means, } 2ab = 2ba$$

or $2ab = 2ab$ (which is true)

a

* $b = b * a$ for all $a, b \in Z^+$ Therefore,

* is commutative. Step 2: Check for Associative.

Consider

* is associative, then

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

(a)

$$* b)^* c = a * (b * c) \text{ LHS} = (a$$

$$* b)^* c = (2ab)^* c$$

=

$$22ab \ c \text{ RHS} = a * (b * c) = a * 2bc$$

=

$$22bc \ a \text{ This implies LHS} \neq \text{RHS}$$

Therefore,

* is not associative binary operation. **(v) On Z^+ , define a**

* **$b = ab$** Step 1: Check for commutative

Consider

* is commutative, then a

$$* b = b * a \text{ Which means, } ab = ba$$

Which is not true

a

$$* b = b * a \text{ for all } a, b \in Z^+ \text{ Therefore,}$$

* is not commutative. Step 2: Check for Associative.

Consider

* is associative, then (a

$$* b)^* c = a * (b * c) \text{ LHS} = (ab)^* c$$

$$= (ab)c$$

$$\text{RHS} = a * (b * c) = a * (bc)$$

$$= abc$$

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

This implies $LHS \neq RHS$

Therefore,

* is not associative. **(vi) On $\mathbb{R} - \{-1\}$, define a**

*** $b = a/(b+1)$** Step 1: Check for commutative

Consider

* is commutative, then a

* $b = b * a$ Which means, $a/(b+1) = b/(a+1)$

Which is not true

Therefore,

* is commutative. Step 2: Check for Associative.

Consider

* is associative, then (a

* $b) * c = a * (b * c)$ LHS = (a

* $b) * c = (a/(b+1)) * c$ a

$$= \frac{b}{c+1}$$

$$= a/(c(b+1))$$

$$RHS = a * (b * c) = a * (b/(c+1))$$

$$= \frac{a}{c+1}$$

$$+1 = a(c+1)/b$$

This implies $LHS \neq RHS$

Therefore,

* is not associative binary operation.

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

3. Consider the binary operation

\wedge on the set $\{1, 2, 3, 4, 5\}$ defined by $a \wedge b = \min \{a, b\}$. Write the operation table of the operation \wedge . **Solution:**

The binary operation

\wedge on the set, say $A = \{1, 2, 3, 4, 5\}$ defined by $a \wedge b = \min \{a, b\}$. the operation table of the operation \wedge is as follows:

\wedge	1	2	3	4	5
1	1	1	1	1	1
2	1	2	2	2	2
3	1	2	3	3	3
4	1	2	3	4	4
5	1	2	3	4	5

4. Consider a binary operation

$*$ on the set $\{1, 2, 3, 4, 5\}$ given by the following multiplication table (Table 1.2).

(i) Compute $(2$

$* 3) * 4$ and $2 * (3 * 4)$ (ii) Is

$*$ commutative? (iii) Compute $(2$

$* 3) * (4 * 5)$. (Hint: use the following table)

Table 1.2

$*$	1	2	3	4	5
1	1	1	1	1	1
2	1	2	1	2	1
3	1	1	3	1	1
4	1	2	1	4	1
5	1	1	1	1	5

Solution:

(i) Compute $(2$

$* 3) * 4$ and $2 * (3 * 4)$ From table: $(2$

$* 3) = 1$ and $(3 * 4) = 1$

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

(2

* 3) * 4 = 1 * 4 = 1 and 2

* (3 * 4) = 2 * 1 = 1 **(ii) Is**

* **commutative?** Consider 2 * 3, we have 2 * 3 = 1 and 3 * 2 = 1

Therefore, * is commutative.

(iii) Compute (2

* 3) * (4 * 5). From table: (2

* 3) = 1 and (4 * 5) = 1 So (2

* 3) * (4 * 5) = 1 * 1 = 1 **5. Let**

*' be the binary operation on the set {1, 2, 3, 4, 5} defined by a *' b = H.C.F. of a and b. Is the operation *' same as the operation * defined in Exercise 4 above? Justify your answer.

Solution: Let A = {1, 2, 3, 4, 5} and a

*' b H.C.F. of a and b. Plot a table values, we have

*'	1	2	3	4	5
1	1	1	1	1	1
2	1	2	1	2	1
3	1	1	3	1	1
4	1	2	1	4	1
5	1	1	1	1	5

Operation

*' same as the operation *. **6. Let**

b = L.C.M. of a and b. Find

* be the binary operation on N given by a *(i) 5

* 7, 20 * 16 (ii) Is

* commutative? (iii) Is

* associative? (iv) Find the identity of

* in N

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

(v) Which elements of \mathbb{N} are invertible for the operation

*? Solution:

(i) 5

* $7 = \text{LCM of } 5 \text{ and } 7 = 35$

* $16 = \text{LCM of } 20 \text{ and } 16 = 80$ (ii) Is

* commutative? a

* $b = \text{L.C.M. of } a \text{ and } b$

* $a = \text{L.C.M. of } b \text{ and } a$

* $b = b * a$ Therefore

* is commutative. (iii) Is

* associative? For a, b, c

$\in \mathbb{N}$
(a

* $b) * c = (\text{L.C.M. of } a \text{ and } b) * c = \text{L.C.M. of } a, b \text{ and } c$

* $(b * c) = a * (\text{L.C.M. of } b \text{ and } c) = \text{L.C.M. of } a, b \text{ and } c$ (a

* $b) * c = a * (b * c)$ Therefore, operation

* associative. (iv) Find the identity of

* in \mathbb{N} Identity of $*$ in $\mathbb{N} = 1$

because $a * 1 = \text{L.C.M. of } a \text{ and } 1 = a$

(v) Which elements of \mathbb{N} are invertible for the operation

? Only the element 1 in \mathbb{N} is invertible for the operation $$ because $1 * 1/1 = 1$

7. Is

* defined on the set $\{1, 2, 3, 4, 5\}$ by $a * b = \text{L.C.M. of } a \text{ and } b$ a binary operation? Justify your answer

Solution:

The operation

* defined on the set $\{1, 2, 3, 4, 5\}$ by $a * b = \text{L.C.M. of } a \text{ and } b$

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

Suppose, $a = 2$ and $b = 3$

$$2 * 3 = \text{L.C.M. of } 2 \text{ and } 3 = 6$$

But 6 does not belong to the set A.

Therefore, given operation $*$ is not a binary operation.

8. Let

$*$ be the binary operation on \mathbb{N} defined by $a * b = \text{H.C.F. of } a \text{ and } b$. Is $*$ commutative? Is $*$ associative? Does there exist identity for this binary operation on \mathbb{N} ? Solution:

The operation

$*$ be the binary operation on \mathbb{N} defined by $a * b = \text{H.C.F. of } a \text{ and } b$ $a * b = \text{H.C.F. of } a \text{ and } b = \text{H.C.F. of } b \text{ and } a$

Therefore, operation $*$ is commutative.

$$\text{Again, } (a * b) * c = (\text{HCF of } a \text{ and } b) * c = \text{HCF of } (\text{HCF of } a \text{ and } b) \text{ and } c = a * (b * c)$$

$$(a * b) * c = a * (b * c)$$

Therefore, the operation is associative.

$$\text{Now, } 1 * a = a * 1 \neq a$$

Therefore, there does not exist any identity element.

9. Let

$*$ be a binary operation on the set \mathbb{Q} of rational numbers as follows: (i) a

$*$ $b = a - b$ (ii) a

$*$ $b = a^2 + b^2$ (iii) a

$*$ $b = a + ab$ (iv) a

$*$ $b = (a - b)^2$ (v) a

$*$ $b = ab/4$ (vi) a

$*$ $b = ab^2$ Find which of the binary operations are commutative and which are associative.

Solution:

(i) a

$*$ $b = a - b$ a

$$* b = a - b = -(b - a) = -b * c \neq b * a \text{ (Not commutative)} \quad (a * b) * c = (a - b) * c = (a - (b - c)) = a - b + c \neq a * (b * c)$$

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

(ii) a

$$* b = a^2 + b^2 a$$

$* b = a^2 + b^2 = b^2 + a^2 = b * a$ (operation is commutative) Check for associative:

$$(a * b) * c = (a^2 + b^2) * c^2 = (a^2 + b^2) + c^2$$

$$a * (b * c) = a * (b^2 + c^2) = a^2 * (b^2 + c^2)^2$$

$(a * b) * c \neq a * (b * c)$ (Not associative)

(iii) a

$$* b = a + ab a$$

$$* b = a + ab = a(1 + b) \quad b * a = b + ba = b(1+a)$$

a

$* b \neq b * a$ The operation $*$ is not commutative

Check for associative:

$$(a * b) * c = (a + ab) * c = (a + ab) + (a + ab)c$$

$$a * (b * c) = a * (b + bc) = a + a(b + bc)$$

$$(a * b) * c \neq a * (b * c)$$

The operation $*$ is not associative

(iv) a

$$* b = (a - b)^2 a$$

$$* b = (a - b)^2 \quad b * a = (b - a)^2$$

a

$* b = b * a$ The operation $*$ is commutative.

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

Check for associative:

$$(a * b) * c = (a - b)^2 * c = ((a - b)^2 - c)^2$$

$$a * (b * c) = a * (b - c)^2 = (a - (b - c)^2)^2$$

$$(a * b) * c \neq a * (b * c)$$

The operation $*$ is not associative

(v) a

$$* \mathbf{b} = \mathbf{ab/4} \quad b * a = ba/2 = ab/2$$

a

$$* b = b * a \quad \text{The operation } * \text{ is commutative.}$$

Check for associative:

$$(a * b) * c = ab/4 * c = abc/16$$

$$a * (b * c) = a * (bc/4) = abc/16$$

$$(a * b) * c = a * (b * c)$$

The operation $*$ is associative.

(vi) a

$$* \mathbf{b} = \mathbf{ab^2} \quad b$$

$$* a = ba^2 \quad a$$

$$* b \neq b * a \quad \text{The operation } * \text{ is not commutative.}$$

Check for associative:

$$(a * b) * c = (ab^2) * c = ab^2 c^2$$

$$a * (b * c) = a * (b c^2) = ab^2 c^4$$

$$(a * b) * c \neq a * (b * c)$$

The operation $*$ is not associative.

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

10. Find which of the operations given above has identity.

Solution: Let I be the identity.

$$(i) a * I = a - I \neq a$$

$$(ii) a * I = a^2 - I^2 \neq a$$

$$(iii) a * I = a + a I \neq a$$

$$(iv) a * I = (a - I)^2 \neq a$$

$$(v) a * I = aI/4 \neq a$$

Which is only possible at $I = 4$ i.e. $a * I = aI/4 = a(4)/4 = a$

$$(vi) a * I = a I^2 \neq a$$

Above identities does not have identity element except (V) at $b = 4$.

11. Let $A = N \times N$ and

$*$ be the binary operation on A defined by (a, b)

$*$ $(c, d) = (a + c, b + d)$ Show that

on A , if any.

$*$ is commutative and associative. Find the identity element for $*$ **Solution:** $A = N \times N$ and $*$ is a binary o

$$(a, b) * (c, d) = (a + c, b + d)$$

$$(c, d) * (a, b) = (c + a, d + b) = (a + c, b + d)$$

The operation $*$ is commutative

$$\text{Again, } ((a, b) * (c, d)) * (e, f) = (a + c, b + d) * (e, f)$$

$$= (a + c + e, b + d + f)$$

$$(a, b) * ((c, d) * (e, f)) = (a, b) * (c+e, e+f) = (a+c+e, b+d+f)$$

$$\Rightarrow ((a, b) * (c, d)) * (e, f) = (a, b) * ((c, d) * (e, f))$$

The operation $*$ is associative.

Let (e, f) be the identity function, then

$$(a, b) * (e, f) = (a + e, b + f)$$

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

For identity function, $a = a + e \Rightarrow e = 0$ and $b = b + f \Rightarrow f = 0$

As zero is not a part of set of natural numbers. So identity function does not exist.

As 0

$\notin \mathbb{N}$, therefore, identity-element does not exist.

12. State whether the following statements are true or false. Justify.

(i) For an arbitrary binary operation $*$ on a set \mathbb{N} , $a * a = a$

$\forall a \in \mathbb{N}$. (ii) If $*$ is a commutative binary operation on \mathbb{N} , then $a * (b * c) = (c * b) * a$

Solution:

(i) Given: $*$ being a binary operation on \mathbb{N} , is defined as $a * a = a$

$\forall a \in \mathbb{N}$ Here operation $*$ is not defined, therefore, the given statement is not true.

(ii) Operation $*$ being a binary operation on \mathbb{N} .

$$c * b = b * c$$

$$(c * b) * a = (b * c) * a = a * (b * c)$$

Thus, $a * (b * c) = (c * b) * a$, therefore the given statement is true.

13. Consider a binary operation

$*$ on \mathbb{N} defined as $a * b = a^3 + b^3$. Choose the correct answer. (A) Is

$*$ both associative and commutative? (B) Is

$*$ commutative but not associative? (C) Is

$*$ associative but not commutative? (D) Is

$*$ neither commutative nor associative? Solution: A binary operation

$*$ on \mathbb{N} defined as $a * b = a^3 + b^3$, Also, a

$* b = a^3 + b^3 = b^3 + a^3 = b * a$ The operation $*$ is commutative.

Again, (a

$$* b) * c = (a^3 + b^3) * c = (a^3 + b^3)^3 + c^3 \quad a * (b * c) = a * (b^3 + c^3) = a^3 + (b^3 + c^3)^3$$

(a

$* b) * c \neq a * (b * c)$ The operation $*$ is not associative.

Therefore, option (B) is correct.

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

1. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be defined as $f(x) = 10x + 7$. Find the function $g : \mathbb{R} \rightarrow \mathbb{R}$ such that $g \circ f = f \circ g = \text{IR}$.

Solution:

Firstly, Find the inverse of f .

Let say, g is inverse of f and

$$y = f(x) = 10x + 7$$

$$y = 10x + 7$$

$$\text{or } x = (y-7)/10$$

$$\text{or } g(y) = (y-7)/10; \text{ where } g : Y \rightarrow N$$

$$\text{Now, } g \circ f = g(f(x)) = g(10x + 7)$$

$$(10x + 7) - 7 =$$

$$10x$$

$$= x$$

$$\text{Again, } f \circ g = f(g(x)) = f((y-7)/10)$$

$$= 10((y-7)/10) + 7$$

$$= y - 7 + 7 = y$$

$$= \text{IR}$$

Since $g \circ f = f \circ g = \text{IR}$. f is invertible, and

$$\text{Inverse of } f \text{ is } x = g(y) = (y-7)/10$$

2. Let $f : W \rightarrow W$ be defined as $f(n) = n - 1$, if n is odd and $f(n) = n + 1$, if n is even. Show that f is invertible. Find the inverse of f . Here, W is the set of all whole numbers.

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

Solution:

$f : W \rightarrow W$ be defined as $f(n) = n - 1$, if n is odd and $f(n) = n + 1$, if n is even.

Function can be defined as:

$$f(n) = \begin{cases} n-1, & \text{if } n \text{ is odd} \\ n+1, & \text{if } n \text{ is even} \end{cases}$$

f is invertible, if f is one-one and onto.

For one-one:

There are 3 cases:

for any n and m two real numbers:

Case 1: n and m : both are odd

$$f(n) = n + 1$$

$$f(m) = m + 1$$

$$\text{If } f(n) = f(m)$$

$$\Rightarrow n + 1 = m + 1$$

$$\Rightarrow n = m$$

Case 2: n and m : both are even

$$f(n) = n - 1$$

$$f(m) = m - 1$$

$$\text{If } f(n) = f(m)$$

$$\Rightarrow n - 1 = m - 1$$

$$\Rightarrow n = m$$

Case 3: n is odd and m is even

$$f(n) = n + 1$$

$$f(m) = m - 1$$

$$\text{If } f(n) = f(m)$$

$$\Rightarrow n + 1 = m - 1$$

$$\Rightarrow m - n = 2 \text{ (not true, because Even - Odd } \neq \text{ Even)}$$

Therefore, f is one-one

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

Check for onto:

$$f(n) = \begin{cases} n-1, & \text{if } n \text{ is odd} \\ n+1, & \text{if } n \text{ is even} \end{cases}$$

Say $f(n) = y$, and y

$\in W$ **Case 1: if $n = \text{odd}$**

$$f(n) = n - 1$$

$$n = y + 1$$

Which show, if n is odd, y is even number.

Case 2: If n is even

$$f(n) = n + 1$$

$$y = n + 1$$

$$\text{or } n = y - 1$$

If n is even, then y is odd.

In any of the cases y and n are whole numbers.

This shows, f is onto.

Again, For inverse of f

$$f^{-1} : y = n - 1$$

$$\text{or } n = y + 1 \text{ and } y = n + 1$$

$$n = y - 1$$

$$f^{-1}(n) = \begin{cases} n-1, & \text{if } n \text{ is odd} \\ n+1, & \text{if } n \text{ is even} \end{cases}$$

Therefore, $f^{-1}(y) = y$. This show inverse of f is f itself.

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

3. If $f : \mathbb{R} \rightarrow \mathbb{R}$ is defined by $f(x) = x^2 - 3x + 2$, find $f(f(x))$.

Solution:

$$\text{Given: } f(x) = x^2 - 3x + 2$$

$$f(f(x)) = f(x^2 - 3x + 2)$$

$$= (x^2 - 3x + 2)^2 - 3(x^2 - 3x + 2) + 2$$

$$= x^4 - 6x^3 + 10x^2 - 3x$$

4. Show that the function $f : \mathbb{R} \rightarrow \{x$

$\in \mathbb{R} : -1 < x < 1\}$ defined by $f(x) = \frac{x}{1+|x|}$ is one and onto function.

$$\frac{x}{1+|x|}, x \in \mathbb{R} \text{ is one}$$

Solution:

The function $f : \mathbb{R} \rightarrow \{x$

$\in \mathbb{R} : -1 < x < 1\}$ defined by $f(x) = \frac{x}{1+|x|}$, $x \in \mathbb{R}$ **For one-one:**

Say x, y

$\in \mathbb{R}$ As per definition of $|x|$;

$$f(x) = \begin{cases} -\frac{x}{1-x}, & x < 0 \\ \frac{x}{1+x}, & x \geq 0 \end{cases}$$

$$\text{So } f(x) = \begin{cases} \frac{x}{1-x}, & x < 0 \\ \frac{x}{1+x}, & x \geq 0 \end{cases}$$

For $x \geq 0$

$$f(x) = \frac{x}{1+x}$$

$$f(y) = \frac{y}{1+y}$$

If $f(x) = f(y)$, then

$$\frac{x}{1+x} = \frac{y}{1+y}$$

$$x(1+y) = y(1+x)$$

$$x = y$$

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

For $x < 0$

$$f(x) = x/(1-x)$$

$$f(y) = y/(1-y)$$

If $f(x) = f(y)$, then

$$x/(1-x) = y/(1-y)$$

$$x(1-y) = y(1-x)$$

$$x = y$$

In both the conditions, $x = y$.

Therefore, f is one-one.

Again for onto:

x

$$f(x) = \frac{x}{1-x} \equiv y$$

For $x < 0$

$$y = f(x) = x/(1-x)$$

$$y(1-x) = x$$

$$\text{or } x(1+y) = y$$

$$\text{or } x = y/(1+y) \dots(1)$$

For $x \geq 0$

$$y = f(x) = x/(1+x)$$

$$y(1+x) = x$$

$$\text{or } x = y/(1-y) \dots(2)$$

Now we have two different values of x from both the case.

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

Since y

$\in \{x \in \mathbb{R} : -1 < x < 1\}$ The value of y lies between -1 to 1 .

If $y = 1$

$x = y/(1-y)$ (not defined)

If $y = -1$

$x = y/(1+y)$ (not defined)

So x is defined for all the values of y , and x

$\in \mathbb{R}$ This shows that, f is onto.

Answer: f is one-one and onto.

5. Show that the function $f : \mathbb{R} \rightarrow \mathbb{R}$ given by $f(x) = x^3$ is injective.

Solution:

The function $f : \mathbb{R} \rightarrow \mathbb{R}$ given by $f(x) = x^3$

Let $x, y \in \mathbb{R}$ such that $f(x) = f(y)$

This implies, $x^3 = y^3$

$x = y$

f is one-one. So f is injective.

6. Give examples of two functions $f : \mathbb{N} \rightarrow \mathbb{Z}$ and $g : \mathbb{Z} \rightarrow \mathbb{Z}$ such that $g \circ f$ is injective but g is not injective.

(Hint : Consider $f(x) = x$ and $g(x) = |x|$)

Solution:

Given: two functions are $f : \mathbb{N} \rightarrow \mathbb{Z}$ and $g : \mathbb{Z} \rightarrow \mathbb{Z}$

Let us say, $f(x) = x$ and $g(x) = x$

$g \circ f = (g \circ f)(x) = f(f(x)) = g(x)$

Here $g \circ f$ is injective but g is not.

Let us take an example to show that g is not injective: Since $g(x) = |x|$

$g(-1) = |-1| = 1$ and $g(1) = |1| = 1$

But $-1 \neq 1$

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

7. Give examples of two functions $f : \mathbb{N} \rightarrow \mathbb{Z}$ and $g : \mathbb{Z} \rightarrow \mathbb{Z}$ such that $g \circ f$ is injective but g is not injective.

int : Consider $f(x)$

$$\{x-1 \text{ if } x > 1 \text{ (H)} = x + 1 \text{ and } g(x) = 1$$

if $x=1$) 1 Solution:

Given: Two functions $f : \mathbb{N} \rightarrow \mathbb{Z}$ and $g : \mathbb{Z} \rightarrow \mathbb{Z}$

Say $f(x) = x + 1$

$$g(x) = \begin{cases} x - 1 & \text{if } x > 1 \\ 1 & \text{if } x = 1 \end{cases}$$

1 = 1 if $x=1$ Check if f is onto:

$f : \mathbb{N} \rightarrow \mathbb{N}$ be $f(x) = x + 1$

say $y = x + 1$

or $x = y - 1$

for $y = 1$, $x = 0$, does not belong to \mathbb{N}

Therefore, f is not onto.

Find $g \circ f$

For $x = 1$; $g \circ f = g(x + 1) = 1$ (since $g(x) = 1$)

For $x > 1$; $g \circ f = g(x + 1) = (x + 1) - 1 = x$ (since $g(x) = x - 1$)

So we have two values for $g \circ f$.

As $g \circ f$ is a natural number, as $y = x$. x is also a natural number. Hence $g \circ f$ is onto.

8. Given a non empty set X , consider $P(X)$ which is the set of all subsets of X .

Define the relation R in $P(X)$ as follows:

For subsets A, B in $P(X)$, $A R B$ if and only if A

$\subset B$. Is R an equivalence relation on $P(X)$? Justify your answer.

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

Solution:

$A \subseteq A \therefore R$ is reflexive.

$A \subseteq B \neq B \subseteq A \therefore R$ is not commutative.

If $A \subseteq B, B \subseteq C$, then $A \subseteq C \therefore R$ is transitive

Therefore, R is not equivalent relation

9. Given a non-empty set X , consider the binary operation $*$: $P(X) \times P(X) \rightarrow P(X)$ given by $A * B = A \cap B \forall A, B$ in $P(X)$, where $P(X)$ is the power set of X . Show that X is the identity element for this operation and X is the only invertible element in $P(X)$ with respect to the operation

* .Solution:

Let T be a non-empty set and $P(T)$ be its power set. Let any two subsets A and B of T .

$$A \cup B \subset T$$

$$\text{So, } A \cup B \in P(T)$$

Therefore, \cup is an binary operation on $P(T)$.

Similarly, if $A, B \in P(T)$ and $A - B \in P(T)$, then the intersection of sets and difference of sets are also binary operation on $P(T)$ and $A \cap T = A = T \cap A$ for every subset A of sets $A \cap T = A = T \cap A$ for all $A \in P(T)$

T is the identity element for intersection on $P(T)$.

10. Find the number of all onto functions from the set $\{1, 2, 3, \dots, n\}$ to itself.

Solution:

Step 1: Compute the total number of one-one functions in the set $\{1, 2, 3\}$

As f is onto, every element of $\{1, 2, 3\}$ will have a unique pre-image

Element Number of possible pairings

1 3

2 2

3 1

Total number of one-one function
 $= 3 \times 2 \times 1$
 $= 6$

Step 2 - Compute the total number of onto functions in the given set
 As f is onto, every element of $\{1, 2, 3, \dots, n\}$ will have a unique pre-image

Element Number of possible pairings

$1 \ n$
 $2 \ n - 1$
 $3 \ n - 2$
 \dots
 \dots
 $n - 1 \ 2$
 $n \ 1$

Total number of one-one function
 $= n \times (n - 1) \times (n - 2) \times \dots \times 2 \times 1$
 $= n!$

Hence, the number of all onto functions from the set $\{1, 2, 3, \dots, n\}$ to itself is $n!$.

11. Let $S = \{a, b, c\}$ and $T = \{1, 2, 3\}$. Find F^{-1} of the following functions F from S to T , if it exists.

(i) $F = \{(a, 3), (b, 2), (c, 1)\}$ (ii) $F = \{(a, 2), (b, 1), (c, 1)\}$

Solution:

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

$$(i) F = \{(a, 3), (b, 2), (c, 1)\}$$

$$F(a) = 3, F(b) = 2 \text{ and } F(c) = 1$$

$$F^{-1}(3) = a, F^{-1}(2) = b \text{ and } F^{-1}(1) = c$$

$$F^{-1} = \{(3, a), (2, b), (1, c)\}$$

$$(ii) F = \{(a, 2), (b, 1), (c, 1)\}$$

Since element b and c have the same image 1 i.e. (b, 1), (c, 1).

Therefore, F is not one-one function.

12. Consider the binary operations

$*$: $\mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ and \circ : $\mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ defined as $a * b = |a - b|$ and $a \circ b = a$,

$\forall a, b \in \mathbb{R}$. Show that $*$ is commutative but not associative, \circ is associative but not commutative. Further

$$\forall a, b, c \in \mathbb{R}, a * (b \circ c) = (a * b) \circ a$$

$* \circ$. [If it is so, we say that the operation $*$ distributes over the operation \circ]. Does \circ distribute over

$*$? Justify your answer. **Solution:**

Step 1: Check for commutative and associative for operation $*$.

$$a * b = |a - b| \text{ and } b * a = |b - a| = |a - b|$$

Operation $*$ is commutative.

$$a * (b * c) = a * |b - c| = |a - (b - c)| = |a - b + c| \text{ and}$$

$$(a * b) * c = |a - b| * c = |a - b - c|$$

$$\text{Therefore, } a * (b * c) \neq (a * b) * c$$

Operation $*$ is associative.

Step 2: Check for commutative and associative for operation \circ .

$$a \circ b = a$$

$$\forall a, b \in \mathbb{R} \text{ and } b \circ a = b \text{ This implies } a \circ b \neq b \circ a$$

Operation \circ is not commutative.

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

Again, $a \circ (b \circ c) = a \circ b = a$ and $(a \circ b) \circ c = a \circ c = a$
 Here $a \circ (b \circ c) = (a \circ b) \circ c$

Operation \circ is associative.

Step 3: Check for the distributive properties

If $*$ is distributive over \circ then $a * (b \circ c) = a * b \circ a * c$

RHS:

$$(a * b) \circ (a * c) = (a - b) \circ (a - c) = |a - b|$$

= LHS

And, $a \circ (b * c) = (a \circ b) * (a \circ c)$

LHS

$$a \circ (b * c) = a \circ (|b - c|) = a$$

RHS $(a \circ b) * (a \circ c) = a * a = |a - a| = 0$

LHS \neq RHS

Hence, operation \circ does not distribute over $*$.

13. Given a non-empty set X, let

$*$: $P(X) \times P(X) \rightarrow P(X)$ be defined as

$A * B = (A - B) \cup (B - A)$, $\forall A, B \in P(X)$. Show that the empty set ϕ is the identity for the operation $*$ and all the elements A of $P(X)$ are invertible with $A^{-1} = A$. (Hint : $(A - \phi) \cup (\phi - A) = A$ and $(A - A) \cup (A - A) = \phi$). Solution:

$x \in P(X)$

$$\phi * A = (\phi - A) \cup (A - \phi) = \phi \cup A = A$$

And

$$A * \phi = (A - \phi) \cup (\phi - A) = A \cup \phi = A$$

ϕ is the identity element for the operation $*$ on $P(X)$.

Also $A * A = (A - A) \cup (A - A)$

$$= \phi \cup \phi = \phi$$

Every element A of $P(X)$ is invertible with $A^{-1} = A$.

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

14. Define a binary operation

* on the set {0, 1, 2, 3, 4, 5} as

$$a * b = \begin{cases} a+b & \text{if } a+b < 6 \\ 6-a & \text{if } a+b \geq 6 \end{cases}$$

Show that zero is the identity for this operation and each element $a \neq 0$ of the set is invertible with $6 - a$ being the inverse of a .

Solution:

Let $X = \{0, 1, 2, 3, 4, 5\}$ and operation * is defined as

$$a * b = \begin{cases} a+b & \text{if } a+b < 6 \\ 6-a & \text{if } a+b \geq 6 \end{cases} \quad \forall a, b \in X$$

Let us say, e is the identity for the operation *, if $a * e = a = e * a$

$$\begin{cases} a+b=0=b+a, & \text{if } a+b < 6 \\ a+b-6=0=b+a-6, & \text{if } a+b \geq 6 \end{cases}$$

That is $a = -b$ or $b = 6 - a$, which shows $a \neq -b$

Since $X = \{0, 1, 2, 3, 4, 5\}$ and $a, b \in X$

Inverse of an element a

$\in X$, $a \neq 0$, and $a^{-1} = 6 - a$.

15. Let $A = \{-1, 0, 1, 2\}$, $B = \{-4, -2, 0, 2\}$ and $f, g : A \rightarrow B$ be functions defined by

$$f(x) = x^2 - x, x \in A$$

and $g(x) = 2|x - \frac{1}{2}| - 1, x \in A$. Are f and g equal? Justify your answer. (Hint: One may note that two functions f and g are called equal functions if $f(a) = g(a)$ for all $a \in A$, are called equal functions).

Solution:

Given functions are: $f(x) = x^2 - x$ and $g(x) = 2|x - \frac{1}{2}| - 1$

At $x = -1$

$$f(-1) = 1 + 1 = 2 \text{ and } g(-1) = 2|-1 - \frac{1}{2}| - 1 = 2$$

At $x = 0$

$$f(0) = 0 \text{ and } g(0) = 0$$

At $x = 1$

$$f(1) = 0 \text{ and } g(1) = 0$$

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

At $x = 2$

$$F(2) = 2 \text{ and } g(2) = 2$$

So we can see that, for each a

$\in A$, $f(a) = g(a)$ This implies f and g are equal functions.

16. Let $A = \{1, 2, 3\}$. Then number of relations containing $(1, 2)$ and $(1, 3)$ which are reflexive and symmetric but not transitive is

(A) 1 (B) 2 (C) 3 (D) 4

Solution:

Option (A) is correct.

As 1 is reflexive and symmetric but not transitive.

17. Let $A = \{1, 2, 3\}$. Then number of equivalence relations containing $(1, 2)$ is

(A) 1 (B) 2 (C) 3 (D) 4

Solution:

Option (B) is correct.

18. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be the Signum Function defined as

$$f(x) = \begin{cases} 1, & x > 0 \\ 0, & x = 0 \\ -1, & x < 0 \end{cases}$$

and $g : \mathbb{R} \rightarrow \mathbb{R}$ be the Greatest Integer Function given by $g(x) = [x]$, where $[x]$ is greatest integer less than or equal to x . Then, does $f \circ g$ and $g \circ f$ coincide in $(0, 1]$?

Solution:

Given:

$f : \mathbb{R} \rightarrow \mathbb{R}$ be the Signum Function defined as

$$f(x) = \begin{cases} 1, & x > 0 \\ 0, & x = 0 \\ -1, & x < 0 \end{cases}$$

and $g : \mathbb{R} \rightarrow \mathbb{R}$ be the Greatest Integer Function given by $g(x) = [x]$, where $[x]$ is

NCERT Solutions for Class 12 Maths Chapter 1 Relations and Functions

greatest integer less than or equal to x .

Now, let say x

$\in (0, 1]$, then $[x] = 1$ if $x = 1$ and

$[x] = 0$ if $0 < x < 1$

Therefore:

$$\begin{aligned} f \circ g(x) &= f(g(x)) = f([x]) \\ &= \begin{cases} f(1), & \text{if } x = 1 \\ f(0), & \text{if } x \in (0, 1) \end{cases} \\ &= \begin{cases} 1, & \text{if } x = 1 \\ 0, & \text{if } x \in (0, 1) \end{cases} \end{aligned}$$

$$\text{Gof}(x) = g(f(x)) = g(1) = [1] = 1$$

For $x > 0$

When x

$\in (0, 1)$, then $f \circ g = 0$ and $\text{gof} = 1$ But $f \circ g(1) \neq \text{gof}(1)$

This shows that, $f \circ g$ and gof do not coincide in $(0, 1]$.

19. Number of binary operations on the set $\{a, b\}$ are

(A) 10 (B) 16 (C) 20 (D) 8

Solution:

Option (B) is correct.

$A = \{a, b\}$ and

$A \times A = \{(a,a), (a,b), (b,b), (b,a)\}$

Number of elements = 4

So, number of subsets = $2^4 = 16$.